



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Anti-Lipopolysaccharide [S69-4] Bulk Size Ab04122-1.1-BT

Isotype and Format: Mouse IgG1, Kappa

Clone Number: S69-4

Alternative Name(s) of Target: LPS; Kdo(2→8)[Kdo(2→4)]Kdo(2→4) Kdo; Kdo₄; branched Kdo₄ tetrasaccharide; *Chlamydomphila psittaci* Lipopolysaccharide tetrasaccharide; 3-deoxy-alpha-d-manno-oct-2-ulopyranosonic acid

UniProt Accession Number of Target Protein:

Published Application(s): SPR, TLC, ELISA, IF

Published Species Reactivity: *Chlamydomphila psittaci*

Immunogen: The original antibody was generated by immunizing BALB/c mice with a synthetic neoglycoconjugate antigen — Kdo(2→8)[Kdo(2→4)]Kdo(2→4) Kdo-BSA (Kdo₄-BSA).

Specificity: This antibody is specific for Kdo₄, a *Chlamydomphila psittaci* Lipopolysaccharide (LPS) tetrasaccharide (3-deoxy-alpha-d-manno-oct-2-ulopyranosonic acid (Kdo)) of the sequence Kdo(2→8)[Kdo(2→4)] Kdo(2→4)Kdo. This antibody specifically binds to the LPS of *Chlamydomphila psittaci* but not to LPS from *Chlamydomphila pneumoniae* or *Chlamydia trachomatis*. Furthermore, it binds exclusively to the LPS and O-deacylated LPS of *Chlamydomphila psittaci*.

Application Notes: This antibody demonstrated high affinity and specificity for structures containing the branched Kdo₄ tetrasaccharide or the 2→4/2→4Kdo₃ trisaccharide, enabling differentiation of *Chlamydomphila psittaci* from other chlamydial species. Furthermore, it shows potential for detecting individual molecular lipopolysaccharide (LPS) species and studying the enzymology of Kdo transferases. Its binding affinity to neoglycoconjugate antigens was assessed using ELISA checkerboard titrations. It displayed comparable affinity for 2→4/2→4Kdo₃-BSA and Kdo₄-BSA and weak binding was observed with 2→4Kdo₂-BSA, 2→8Kdo₂-BSA, and 2→8/2→4Kdo₃-BSA, but only at high antibody and antigen concentrations. This antibody was also tested against BSA glycoconjugates containing 4'-monophosphoryl derivatives of deacylated *C. psittaci* LPS. It demonstrated similar reactivity with 2→4/2→4Kdo₃-GlcNAc₂-4P-BSA and Kdo₄-GlcNAc₂-4P-BSA. Further experiments confirmed its binding to isolated LPS and whole bacteria. Inhibition assays using free oligosaccharides as inhibitors in ELISA demonstrated that the antibody achieved 50% inhibition with 2→4/2→4Kdo₃-allyl, Kdo₄-allyl, and Kdo₄-GlcNAc₂-P₂ at concentrations of 4.4 μM, 1.7 μM, and 2.1 μM, respectively. An immunofluorescence (IF) assay on infected L929 and HL cells confirmed the antibody's recognition of a consistent epitope in *C. psittaci* inclusions, while remaining negative for *Chlamydomphila pneumoniae* and *Chlamydia trachomatis*. Additional IF testing using various chlamydial strains confirmed the antibody's specificity, as all four tested *C. psittaci* strains displayed positive staining. Surface plasmon resonance (SPR) binding studies were conducted to determine the

interaction between the original antibody format (mouse IgG1) with various ligands; the dissociation constants (K_d) for the immobilized antibody were measured for Kdo₄-GlcNAc₂-P₂, 2→4/2→4Kdo₃-GlcNAc₂-P₂, and 2→8/2→4Kdo₃-GlcNAc₂-P₂, resulting in K_d values of 10 μM, 20 μM, and 200 μM, respectively. When the immobilized scFv version of the antibody was used, the binding constants at 150 mM NaCl were 10 μM, 100 μM, and 800 μM for Kdo₄-GlcNAc₂-P₂, 2→4/2→4Kdo₃-GlcNAc₂-P₂, and 2→8/2→4Kdo₃-GlcNAc₂-P₂, respectively. The SPR studies revealed that this antibody exhibits a higher affinity for Kdo₄-GlcNAc₂-P₂ compared to 2→4/2→4Kdo₃-GlcNAc₂-P₂, and significantly reduced affinity for the Kdo-trisaccharide containing 2→8-linked terminal Kdo (Müller-Loennies et al., 2006; PMID: 16282606).

Antibody First Published in: Müller-Loennies et al. A monoclonal antibody against a carbohydrate epitope in lipopolysaccharide differentiates *Chlamydomphila psittaci* from *Chlamydomphila pecorum*, *Chlamydomphila pneumoniae*, and *Chlamydia trachomatis* *Glycobiology*. 2006 Mar;16(3):184-96. doi: 10.1093/glycob/cwj055 PMID:16282606

Note on publication: The original publication discusses the development of a monoclonal antibody that can differentiate *Chlamydomphila psittaci* from other *Chlamydomphila* species and *Chlamydia trachomatis* based on a specific carbohydrate epitope in the lipopolysaccharide.

Product Form

Size: 1 mg Purified antibody in bulk size.

Purification: Protein A affinity purified

Supplied In: PBS only.

Storage Recommendation: Store at 4°C for up to 3 months. Note, this antibody is provided without added preservatives, it is therefore recommended this antibody be handled under sterile conditions. For longer storage, aliquot and store at -20°C.

Concentration: 1 mg/ml.

Important note - This product is for research use only. It is not intended for use in therapeutic or diagnostic procedures for humans or animals.