



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## Anti-RANK-L [3K7F5] Standard Size Ab04140-23.0

**Isotype and Format:** Rabbit IgG, Kappa

**Clone Number:** 3K7F5

**Alternative Name(s) of Target:** CD254; TRANCE; RANKL; PAE30; ODF; OPGL; Tumor necrosis factor ligand superfamily member 11; Osteoclast differentiation factor; Osteoprotegerin ligand; Receptor activator of nuclear factor kappa-B ligand; TNF-related activation-induced cytokine

**UniProt Accession Number of Target Protein:** O14788

**Published Application(s):** in vivo

**Published Species Reactivity:** Human

**Immunogen:** The original antibody was generated by panning a phage display library against human sRANKL.

**Specificity:** The antibody is specific for human-RANK-L. RANKL is a tumor necrosis factor related protein that binds to, and activates, the TNF receptor related protein Receptor Activator of NF-κB (RANK). RANKL acts as the key cytokine that regulates osteoclast differentiation and activation during normal bone remodeling and during disease

**Application Notes:** The affinity of the original format of the antibody is  $0.17 \times 10^{-12}$  nM, as measured by PAE technology. RAW264.7, a murine macrophage cell line that can be induced by RANKL to differentiate into osteoclast-like cells, was used in a TRAP test. In the TRAP test, the original antibody or the Fab fragment were able to neutralize the stimulating effect of RANKL on osteoclast formation from a RAW 264.7 culture. Further, the addition of the original format of the antibody or the Fab fragment inhibited osteoclast differentiation of the RAW cell in a dose-dependent manner. The affinity of the original format and the Fab fragment of the antibody was measured by the Scatchard method (affinity = 8.7 and 6.5 pM). The original antibody and the Fab fragment could significantly increase the bone densities and bone calcium content in a dose-dependent manner, in in vivo experiments (WO2011116527A1).

**Antibody First Published in:** PMID:

**Note on publication:**

### Product Form

**Size:** 100 µg Purified antibody.

**Purification:** Protein A affinity purified

**Supplied In:** PBS with 0.02% Proclin 300.

**Storage Recommendation:** Store at 4°C for up to 3 months. For longer storage, aliquot and store at -

20°C.

**Concentration:** 1 mg/ml.

Important note - This product is for research use only. It is not intended for use in therapeutic or diagnostic procedures for humans or animals.