



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Anti-domoic acid [2H12] Bulk Size Ab04145-1.1-BT

Isotype and Format: Mouse IgG1, Kappa

Clone Number: 2H12

Alternative Name(s) of Target: DA; (2S,3S,4S)-4-[(2Z,4E,6R)-6-Carboxyhepta-2,4-dien-2-yl]-3-(carboxymethyl)pyrrolidine-2-carboxylic acid

UniProt Accession Number of Target Protein:

Published Application(s): ELISA

Published Species Reactivity: Specie independent

Immunogen:

Specificity: The antibody is specific for domoic acid. Domoic acid is a kainic acid-type neurotoxin that causes amnesic shellfish poisoning. It is produced by algae and accumulates in shellfish, sardines, and anchovies. When sea lions, otters, cetaceans, humans, and other predators eat contaminated animals, poisoning may result.

Application Notes: The specificity of the original format and the scFv format of the antibody was confirmed by ELISA analysis (IC₅₀ = 1 µg/ml and 245 ng/ml respectively and K_d = 0.8 µM for the scFv fragment) (Hu et al., 2005; PMID: 16019098). The scFv was employed for the development of a novel immunosensor for in situ domoic acid screening, where the fragment was immobilized on mesoporous silica. Domoic acid binding analysis found that the adsorbed scFv bound domoic acid in a concentration-dependent manner (Hu et al., 2007; PMID: 17005419). The scFv antibody fragment was engineered with cysteine-containing linkers of two different lengths. The domoic acid detection ability of the covalently immobilised antibody fragments was compared with that of scFvs adsorbed non-covalently onto polystyrene plates. The IC₅₀ for DA of the unmodified scFv in solution was previously reported as 245 ng/mL whereas in this study, the scFv, 2H12 scFv-cys I and 2H12 scFv-cys II proteins adsorbed onto standard Maxisorp plates were calculated to have IC₅₀s for DA of 540 ng/mL, 1126 ng/mL and 1482 ng/mL, respectively, as measured by Competitive ELISA (Hortigüela et al., 2013; PMID: 23493076).

Antibody First Published in: Hu et al. Cloning, expression and characterisation of a single-chain Fv antibody fragment against domoic acid in *Escherichia coli* J Biotechnol. 2005 Oct 17;120(1):38-45. doi: 10.1016/j.jbiotec.2005.05.018. Epub 2005 Jul 12. [PMID:16019098](#)

Note on publication: The original paper describes the cloning, expression and characterization of a single chain Fv antibody from a mouse hybridoma cell line.

Product Form

Size: 1 mg Purified antibody in bulk size.

Purification: Protein A affinity purified

Supplied In: PBS only.

Storage Recommendation: Store at 4°C for up to 3 months. Note, this antibody is provided without added preservatives, it is therefore recommended this antibody be handled under sterile conditions. For longer storage, aliquot and store at -20°C.

Concentration: 1 mg/ml.

Important note - This product is for research use only. It is not intended for use in therapeutic or diagnostic procedures for humans or animals.