



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## Anti-domoic acid [DA24cB7] Standard Size Ab04146-30.11

This is an scFv fragment with a His tag.

This is a reformatted sheep scFv antibody, based on the original sheep scFv format, created for improved compatibility with existing reagents, assays and techniques.

**Isotype and Format:** scFv fragment (His), ScFv

**Clone Number:** DA24cB7

**Alternative Name(s) of Target:** DA; (2S,3S,4S)-4-[(2Z,4E,6R)-6-Carboxyhepta-2,4-dien-2-yl]-3-(carboxymethyl)pyrrolidine-2-carboxylic acid

**UniProt Accession Number of Target Protein:**

**Published Application(s):** ELISA

**Published Species Reactivity:** Specie independent

**Immunogen:** The original antibody was originated from a scFv phage library obtained from a sheep hyperimmunised with DA-BSA conjugate. It was isolated by panning against DA.

**Specificity:** The antibody is specific for domoic acid. Domoic acid is a kainic acid-type neurotoxin that causes amnesic shellfish poisoning. It is produced by algae and accumulates in shellfish, sardines, and anchovies. When sea lions, otters, cetaceans, humans, and other predators eat contaminated animals, poisoning may result.

**Application Notes:** The specificity of the original format of the antibody was confirmed by competitive ELISA using plates coated with DA-OVA (150 of 1.21 ng/mL (3.9 nM)). The assay was used for the determination of DA levels in shellfish extracts (Shaw et al., 2008; PMID: 18384202).

**Antibody First Published in:** Shaw et al. Development of a high-affinity anti-domoic acid sheep scFv and its use in detection of the toxin in shellfish Anal Chem. 2008 May 1;80(9):3205-12. doi: 10.1021/ac7024199. Epub 2008 Apr 3. [PMID:18384202](#)

**Note on publication:** The paper describes the generation and characterization of the antibody.

### Product Form

**Size:** 100 µg Purified antibody.

**Purification:** Purified by Immobilized Metal Affinity Chromatography

**Supplied In:** PBS with 0.02% Proclin 300.

**Storage Recommendation:** Store at 4°C for up to 3 months. For longer storage, aliquot and store at -

20°C.

**Concentration:** 1 mg/ml.

Important note - This product is for research use only. It is not intended for use in therapeutic or diagnostic procedures for humans or animals.