

# Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

## Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

#### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in





## Anti-Giantin [TA10] Standard Size Ab04207-30.11

This is an scFv fragment with a His tag.

This is a reformatted human scFv antibody, based on the original human scFv format, created for improved compatibility with existing reagents, assays and techniques.

**Isotype and Format:** scFv fragment (His), ScFv

**Clone Number: TA10** 

Alternative Name(s) of Target: Golgin subfamily B member 1; 372 kDa Golgi complex-associated

protein; GCP372; Macrogolgin

**UniProt Accession Number of Target Protein:** 

Published Application(s): in vivo, WB, IF
Published Species Reactivity: Rat, Human

**Immunogen:** The original antibody was generated via phage display from the Griffin.1 library by screening against Rat liver Golgi (RLG) complexes derived from rat liver.

**Specificity:** This antibody is specific for a 79 amino acid long region between amino acids 1462 to 1540 of the full human sequence (corresponding to amino acids 1399 to 1478 in the rat homologue) in Giantin, a Golgi matrix protein.

**Application Notes:** The original format of this antibody (scFv) was successfully used in a Western blot and for *in vitro* immunofluorescence straining of Giantin in HeLa cells. siRNA knock-down of Giantin reduced TA10 staining, confirming its specificity. Furthermore, an *in vivo* expressed TA10-GFP construct successfully stained HeLa cells (Nizak et al., 2003; PMID: 14617357). The scFv-Fc fusion version of this antibody — hTA10 — detected Giantin when cells were fixed by paraformaldehyde (Moutel et al., 2009; PMID: 19245715). Precise characterization of the TA10 target was achieved; a region of 79 amino acids, representing less than 2.5% of the entire protein, was determined as its binding site (Vielemeyer et al., 2010; PMID: 20727208).

**Antibody First Published in:** Nizak et al. Recombinant antibodies against subcellular fractions used to track endogenous Golgi protein dynamics in vivo Traffic. 2003 Nov;4(11):739-53. doi: 10.1034/j.1600-0854.2003.00132.x PMID:14617357

**Note on publication:** The original publication demonstrates the use of recombinant antibodies selected through phage display against subcellular fractions.

#### **Product Form**

**Size:** 100 μg Purified antibody.

Purification: Purified by Immobilized Metal Affinity Chromatography

**Supplied In:** PBS with 0.02% Proclin 300.

Storage Recommendation: Store at 4°C for up to 3 months. For longer storage, aliquot and store at -

20°C.

**Concentration:** 1 mg/ml.

Important note – This product is for research use only. It is not intended for use in therapeutic or diagnostic procedures for humans or animals.