



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Anti-Collagenase type IV [3G11] Standard Size Ab04210-2.3

This antibody was created using our proprietary Fc Silent™ engineered Fc domain containing key point mutations that abrogate binding to Fc gamma receptors.

This is a reformatted mouse IgG2a Fc Silent™ antibody, based on the original mouse IgG1 format, created for improved compatibility with existing reagents, assays and techniques.

Isotype and Format: Mouse IgG2a, [Fc Silent™](#), Kappa

Clone Number: 3G11

Alternative Name(s) of Target: MMP2; MMP-2; MMP 2; Matrix metalloproteinase 2; 72 kDa type IV collagenase; 72 kDa gelatinase; Gelatinase A

UniProt Accession Number of Target Protein: P33434

Published Application(s): Immunoscintigraphy, in vivo, MTT assay, ELISA, IF, IHC

Published Species Reactivity: Mouse

Immunogen: The original antibody was generated by immunizing BALB/c mice with type IV collagenase.

Specificity: This antibody is specific for type IV collagenase.

Application Notes: The immunoreactivity of the original format of this antibody (mouse IgG1, κ) was measured via ELISA and was shown to decrease by approximately 10%-20% after cold iodination with ¹³¹I-iodide. The (¹³¹I)-labeled version of this antibody was demonstrated to be stable *in vitro* for almost 72 h. The blood clearance pattern of (¹²⁵I)-labeled 3G11 was defined as a two-compartment model, with a 7.2 h $T_{1/2\alpha}$ and a 345.2 h $T_{1/2\beta}$, and clear scintigraphic images of human lung carcinoma PG were obtained using (¹²⁵I)-labeled 3G11 at 72 h, which further improved at 120 h (Dai et al., 2003; PMID: 14693044). Lidamycin (LDM) was conjugated to the Fab version of this antibody, and the resulting immunoconjugate maintained most of its immunoreactivity to both type IV collagenase and mouse hepatoma 22 cells, as shown by ELISA, and exhibited more potent cytotoxicity to hepatoma 22 cells than free LDM by MTT assay. Furthermore, when administered intravenously, the Fab'-LDM conjugate was proven to be more effective against the growth of subcutaneously transplanted hepatoma 22 in mice than free LDM in two experiment settings, increasing mice survival time (Fengqiang et al., 2004; PMID: 15382678). The scFv version of this antibody was constructed and used in various configurations for IF and *in vivo* imaging and therapeutic applications (Zhong et al., 2010; PMID: 20303650) (Kong et al., 2015; PMID: 25824464).

Antibody First Published in: Dai et al. [Immunoscintigraphy of anti-type IV collagenase monoclonal antibody in nude mice bearing human lung cancer xenograft] *Ai Zheng*. 2003 Dec;22(12):1243-8.

[PMID:14693044](#)

Note on publication: The original publication focuses on evaluating the tumor-specific distribution of the anti-type IV collagenase monoclonal antibody (mAb) 3G11 through radio imaging in tumor-bearing nude mice.

Product Form

Size: 100 µg Purified antibody.

Purification: Protein A affinity purified

Supplied In: PBS with 0.02% Proclin 300.

Storage Recommendation: Store at 4°C for up to 3 months. For longer storage, aliquot and store at -20°C.

Concentration: 1 mg/ml.

Important note – This product is for research use only. It is not intended for use in therapeutic or diagnostic procedures for humans or animals.