



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Anti-IL-17 [Antibody 7] Standard Size Ab04237-10.0

Isotype and Format: Human IgG1, Lambda

Clone Number: Antibody 7

Alternative Name(s) of Target: IL17; IL-17A; IL17A; CTLA8; Interleukin-17A; IL-17A; Cytotoxic T-lymphocyte-associated antigen 8; CTLA-8; IL17F; IL-17F; Interleukin-17F; IL-17F; Cytokine ML-1

UniProt Accession Number of Target Protein:

Published Application(s): crystallization, functional assays, in vivo, SPR, ELISA

Published Species Reactivity: Dog, Human, Cynomolgus Monkey

Immunogen: The original antibody was produced by immunizing Balb/c mice with purified recombinant human IL-17.

Specificity: This antibody is specific for IL-17; it binds residues 74 (Leu), 85 (Tyr), 86 (His), and 87 (Met) of the mature IL-17 sequence. In humans, it has been shown to bind both IL-17A and IL-17F.

Application Notes: The IC_{50} values for this antibody as measured in an HT1080 assay in response to 1 nM human or cynomolgus monkey (cyno) IL-17A were 0.8 ± 0.2 nM and 4.8 ± 1.6 nM, respectively. Partial inhibition was also observed with IL-17F. Further assays showed cross-reactivity with dog IL-17A. This antibody neutralized the synergistic IL-6 response as measured in an altered human HT1080 assay with an IC_{50} value of 0.24 ± 0.05 nM. $pA_2 = 9.23$ and $K_d = 663$ pM were measured when testing the affinity of this antibody for human IL-17A. Further kinetic analysis on recombinant human IL-17A (*E. coli*), cyno IL-17A (*E. coli*), and human IL-17A-Flag-His10 (HEK) yielded a 1:1 Langmuir K_D and Bivalent Analyte K_D of 0.195 nM, 0.512 nM, and 0.171 nM, and 2.098 nM, 1.340 nM, and 0.41-1.88 nM, appropriately. The biological activity of this antibody was tested extensively in the air pouch of Balb/c; it effectively inhibited IL-17A-induced IL-6 release and the influx of white blood cells in the air pouch, as reflected by low ID_{50} values. This antibody exhibited inhibitory effects on the IL-17A/TNF α synergized responses in post-mortem cartilage explants and synovial fibroblasts with the potential to modulate inflammatory and destructive responses induced by the synergistic action of IL-17A and TNF α . X-ray crystallography of the IL-17A bound Fab version of this antibody indicated that it binds residues 74 (Leu), 85 (Tyr), 86 (His), and 87 (Met) of the mature sequence (US8865166).

Antibody First Published in: [PMID:](#)

Note on publication:

Product Form

Size:

100 µg Purified antibody.

Purification: Protein A affinity purified

Supplied In: PBS with 0.02% Proclin 300.

Storage Recommendation: Store at 4°C for up to 3 months. For longer storage, aliquot and store at -20°C.

Concentration: 1 mg/ml.

Important note - This product is for research use only. It is not intended for use in therapeutic or diagnostic procedures for humans or animals.