



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Anti-EGFR [7A7] Bulk size M bAb0506-BS

This antibody is a knob-into-hole bispecific based on murine IgG2a with an Fc Silent™ mutation, featuring one arm with an anti-mouse CD3e scFv based on the clone 145-2C11.

This is a bispecific, knob-into-hole heterodimerized antibody, with one arm directed against mouse EGFR (Fab) and one arm directed against mouse CD3e (scFv).

Isotype and Format: Mouse IgG2a mCD3e KIH bispecific, [Fc Silent™](#), Kappa

Clone Number: 7A7

Alternative Name(s) of Target: ERBB; ERBB1; HER1; PIG61; mENA; avian erythroblastic leukemia viral (v-erb-b) oncogene homolog; cell growth inhibiting protein 40; cell proliferation-inducing protein 61; proto-oncogene c-ErbB-1; receptor tyrosine-protein kinase erbB-1; epidermal growth f

UniProt Accession Number of Target Protein: P00533

Published Application(s): in vivo, FC, IF, IHC

Published Species Reactivity: Human, Mouse

Immunogen: Recombinant extracellular domain of murine EGFR (rECD-mEGFR).

Specificity: Recognizes the extracellular domain of EGFR.

Application Notes: The antibody was generated in order to establish a mouse model for in vivo pre-clinical evaluation of therapies targeting EGFR. The antibody was able to recognize the murine and human EGFR expressed on 3LL or A431 tumor cell lines by flow cytometry. Moreover, the antibody could immunoprecipitate EGFR of 3LL cell lysate, it was used in immunohistochemistry on mouse skin tissue and mammary adenocarcinoma, and it was able to detect EGFR in Western blots (Garrido et al., 2004; PMID: 15312307). The antibody inhibited EGF-induced EGFR signaling in D122 tumor cells by reducing EGFR phosphorylation as well as the phosphorylation of molecules involved in downstream EGFR signaling; the antibody showed strong anti-metastatic activity on D122 tumors, for which the contribution of T cells is essential (Garrido et al., 2007). The crystal structure of the Fab fragment of the antibody was determined. The specificity of the antibody to extracellular EGFR was confirmed by ELISA, while competition ELISA showed the antibody does not compete with Nimotuzumab. The antibody recognized a binding site located on the EGFR extracellular domain that is not accessible in its most stable conformations, but that becomes exposed upon treatment, such as with tyrosine kinase inhibitor (Talavera et al., 2011; PMID: 21592580). Talavera et al. observed the antibody recognition is strictly dependent on the experimental setup, and results can vary significantly, depending on seemingly minor experimental variations. In this regard, He et al. claimed to be unable to reproduce previous results obtained with the antibody. In their experiments the

antibody did not detect EGFR expression by immunofluorescence or Western blot. Flow cytometry experiments showed surface expression of EGFR in A431 cells, weak staining in 3T3-L1 cells, and no staining on the surface of HPV38 cells. The antibody failed to detect EGFR on the plasma membrane of HPV38 tumour sections and in vivo administration of the antibody in an EGFR-expressing HPV38 tumour model did not have any impact on tumour regression or animal survival (He et al., 2018; PMID: 29552307). However, in 2021 Skibbe and coworker successfully used the antibody to treat mice with established colitis-driven CRCs. In particular, the antibody bound to EGFR by ELISA experiments, in vivo experiments showed the antibody significantly lowered the frequency of large tumors in mice and the antibody induced a significant downregulation of PD-L1 in highly glycolytic CRC cells (Skibbe et al., 2021; PMID: 34830971). Absolute antibody wanted to report these discrepancies in the results of different groups, suggesting that optimization of experimental conditions is crucial for the effectiveness of the antibody.

Antibody First Published in: Garrido G, Sanchez B, Rodriguez HM, Lorenzano P, Alonso D, Fernandez LE. 7A7 MAb: a new tool for the pre-clinical evaluation of EGFR-based therapies. Hybrid Hybridomics. 2004 Jun;23(3):168-75. [PMID:15312307](#)

Note on publication: Describes the generation and characterization of an anti-murine EGFR extracellular domain monoclonal antibody (7A7 MAb) (IgG1).

Product Form

Size: 1 mg Purified antibody in bulk size.

Purification: Protein A affinity purified

Supplied In: PBS only.

Storage Recommendation: Store at 4°C for up to 3 months. Note, this antibody is provided without added preservatives, it is therefore recommended this antibody be handled under sterile conditions. For longer storage, aliquot and store at -20°C.

Concentration: See vial label

Important note – This product is for research use only. It is not intended for use in therapeutic or diagnostic procedures for humans or animals.