

# Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



## Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

## Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

## SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in





## MAP3K14 & CHUK Protein Protein Interaction Antibody Pair

Catalog #: DI0605 規格:[1 Set]

#### List All

#### **Specification**

#### **Product Description:**

This protein protein interaction antibody pair set comes with two antibodies to detect the protein-protein interaction, one against the MAP3K14 protein, and the other against the CHUK protein for use in *in* situ Proximity Ligation Assay. See Publication Reference below.

### **Application Image**


In situ Proximity Ligation Assay (Cell)




1. Incubate with target primary antibodies



2. Add PLA probes PLUS and MINUS



3. Hybridize connector oligos



4. Ligation to form a complete DNA circle



5. Rolling circle amplification




6 Add fluorescent probes to reveal interaction

Reactivity:

Human

Quality Control Protein protein interaction immunofluorescence result.





Representative image of Proximity Ligation Assay of protein-protein interactions between MAP3K14 and CHUK. HeLa cells were stained with anti-MAP3K14 rabbit purified polyclonal antibody 1:1200 and anti-CHUK mouse monoclonal antibody 1:50. Each red dot represents the detection of protein-protein interaction complex. The images were analyzed using an optimized freeware (BlobFinder) download from The Centre for Image Analysis at Uppsala University.

#### Supplied **Product:**

Antibody pair set content:

- 1. MAP3K14 rabbit purified polyclonal antibody (20 ug)
- 2. CHUK mouse monoclonal antibody (40 ug)

\*Reagents are sufficient for at least 30-50 assays using recommended protocols.

**Storage** Instruction:

Store reagents of the antibody pair set at -20°C or lower. Please aliquot to avoid repeated freeze thaw cycle. Reagents should be returned to -

20°C storage immediately after use.

MSDS:

<u>Marian</u> Download

#### **Publication Reference**

 An analysis of protein-protein interactions in cross-talk pathways reveals CRKL as a novel prognostic marker in hepatocellular carcinoma.
Liu CH, Chen TC, Chau GY, Jan YH, Chen CH, Hsu CN, Lin KT, Juang YL, Lu PJ, Cheng HC, Chen MH, Chang CF, Ting YS, Kao CY, Hsiao M, Huang CY. Mol Cell

Proteomics. 2013 Feb 8. [Epub ahead of print]

#### **Applications**

In situ Proximity Ligation Assay (Cell)

CHUK MAP3K14

**Gene Information** 

Entrez GeneID: 9020

Gene Name: MAP3K14

Gene Alias: FTDCR1B, HS, HSNIK, NIK

Gene mitogen-activated protein kinase kinase kinase 14

**Description:** 

Omim ID: 604655

Gene Ontology: Hyperlink

Gene Summary: This gene encodes mitogen-activated protein kinase kinase kinase 14,

which is a serine/threonine protein-kinase. This kinase binds to TRAF2 and stimulates NF-kappaB activity. It shares sequence similarity with several other MAPKK kinases. It participates in an NF-kappaB-inducing signalling cascade common to receptors of the tumour-necrosis/nervegrowth factor (TNF/NGF) family and to the interleukin-1 type-I receptor.

[provided by RefSeq

Other serine/threonine protein-kinase

**Designations:** 

**Gene Information** 

Entrez GeneID: 1147

Gene Name: CHUK

Gene Alias: IKBKA,IKK-alpha,IKK1,IKKA,NFKBIKA,TCF16

Gene conserved helix-loop-helix ubiquitous kinase

**Description:** 

Omim ID: 600664

Gene Ontology: Hyperlink

Gene Summary: This gene encodes a member of the serine/threonine protein kinase

family. The encoded protein, a component of a cytokine-activated protein complex that is an inhibitor of the essential transcription factor NF-kappa-B complex, phosphorylates sites that trigger the degradation of the inhibitor via the ubiquination pathway, thereby activating the

transcription factor. [provided by RefSeq

Page 2 of 3 2016/5/20

Other I-kappa-B kinase 1,I-kappa-B kinase-alpha,IKK-a kinase,IkB kinase alpha subunit, Nuclear factor NFkappaB inhibitor kinase Designations: alpha,OTTHUMP0000020273,conserved helix-loop ubiquitous kinase Interactome MDM2 CSNK2A1 FANCA RAF1 ESR1 MAP3K8 PRKCQ DVL<sub>2</sub> FRAP CDKN1B PRKDC APPL1 TRPC4AP ARRB1 PRKCB1 TNFAIP3 MAP2K4 IRS1 EEF1A2 MAP3K1 PRKCZ CDC37 FOXO3 ATXN1 HSP90AB1 MAP3K7IP3 RAK1 HSP90AA1 CHU MAP3K11 CSF2RA MAP3K5 TNFRSE1A BIRC3 CASP8 CREBBP BIRC4 NRIP1 TRAF3IP2 CALM1 EIF2AK2 HSPA1A DNAJA1 TUBA3C TUBB4 RUVBL1 ACTG HSPA9 FKBP5 CTNNB1 CALM3 RUVBL2 IKBKG CAD MAP3K7 TBK1 VISA MAP3K14 NFKB2 TUBB2B HSPA1L PARP1 KBKAP NFKBIB TNFRSF11A HSPA5 MAP3K3 **EPRS** RELA NCOA3 MYH10 TRAF6 NFKB1 MAP3K7IP1 STAP2 RPS27A PRIM1 PEBP1 TGFBR1

服務條款 | 隱私權政策 | 著作及商標 | 網站地圖

©2016 亞諾法生技股份有限公司 Abnova Corporation. 版權所有.

Page 3 of 3 2016/5/20