



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

XPC FISH Probe

Catalog # : FA0088

規格 : [200 uL]

List All

Specification

Product Description:	Made to order FISH probes for identification of gene amplification using Fluorescent In Situ Hybridization Technique. (Technology)
Supplied Product:	DAPI Counterstain (1500 ng/mL) 250 uL
Storage Instruction:	Store at 4°C in the dark.
Origin:	Human
Source:	Genomic DNA
Notice:	We strongly recommend the customer to use FFPE FISH PreTreatment Kit 1 (Catalog #: KA2375 or KA2691) for the pretreatment of Formalin-Fixed Paraffin-Embedded (FFPE) tissue sections.
Regulation Status:	For research use only (RUO)

Application Image

Fluorescent In Situ Hybridization (Cell)

Applications

Fluorescent In Situ Hybridization (Cell)

 [Protocol Download](#)

Gene Information

Entrez GeneID: [7508](#)

Gene Name: XPC

Gene Alias: XP3,XPCC

Gene Description: xeroderma pigmentosum, complementation group C

Omim ID: [278720](#)

Gene Ontology: [Hyperlink](#)

Gene Summary: This gene encodes a component of the nucleotide excision repair (NER) pathway. There are multiple components involved in the NER pathway, including Xeroderma pigmentosum (XP) A-G and V, Cockayne syndrome (CS) A and B, and trichothiodystrophy (TTD) group A, etc. This component, XPC, plays an important role in the early steps of global genome NER, especially in damage recognition, open complex formation, and repair protein complex formation. Mutations in this gene or some other NER components result in Xeroderma pigmentosum, a rare autosomal recessive disorder characterized by increased sensitivity to sunlight with the development of carcinomas at an early age. Alternatively spliced transcript variants have been found for this gene. [provided by RefSeq]

Other xeroderma pigmentosum group C protein

Designations:

Gene Pathway

[Nucleotide excision repair](#)

Related Disease

[Adenocarcinoma](#) [Adenoma](#) [Ataxia telangiectasia](#) [Azoospermia](#) [Barrett Esophagus](#)
[Bone Neoplasms](#) [Brain Neoplasms](#) [Breast cancer](#) [Breast Diseases](#) [Breast Neoplasms](#)
[Carcinoma](#) [Carcinoma, Basal Cell](#) [Carcinoma, Hepatocellular](#)
[Carcinoma, Non-Small-Cell Lung](#) [Carcinoma, Renal Cell](#) [Carcinoma, Squamous Cell](#)
[Carcinoma, Transitional Cell](#) [Cell Transformation, Neoplastic](#) [Chromosome Aberrations](#)

... see more

[服務條款](#) | [隱私權政策](#) | [著作及商標](#) | [網站地圖](#)

©2017 亞諾法生技股份有限公司 Abnova Corporation. 版權所有.