



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

LMO2 FISH Probe

Catalog # : FA0276

規格 : [200 uL]

List All

Specification

Product Description:	Made to order FISH probes for identification of gene amplification using Fluorescent In Situ Hybridization Technique. (Technology)
Supplied Product:	DAPI Counterstain (1500 ng/mL) 250 uL
Storage Instruction:	Store at 4°C in the dark.
Origin:	Human
Source:	Genomic DNA
Notice:	We strongly recommend the customer to use FFPE FISH PreTreatment Kit 1 (Catalog #: KA2375 or KA2691) for the pretreatment of Formalin-Fixed Paraffin-Embedded (FFPE) tissue sections.
Regulation Status:	For research use only (RUO)

Application Image

Fluorescent In Situ Hybridization (Cell)

Applications

Fluorescent In Situ Hybridization (Cell)

 [Protocol Download](#)

Gene Information

Entrez GeneID: [4005](#)

Gene Name: LMO2

Gene Alias: RBTN2,RBTNL1,RHOM2,TTG2

Gene Description: LIM domain only 2 (rhombotin-like 1)

Omim ID: [180385](#)

Gene Ontology: [Hyperlink](#)

Gene Summary: LMO2 encodes a cysteine-rich, two LIM-domain protein that is required for yolk sac erythropoiesis. The LMO2 protein has a central and crucial role in hematopoietic development and is highly conserved. The LMO2 transcription start site is located approximately 25 kb downstream from the 11p13 T-cell translocation cluster (11p13 ttc), where a number T-cell acute lymphoblastic leukemia-specific translocations occur. Alternative splicing results in multiple transcript variants encoding different isoforms

Other Designations: LIM domain only 2,T-cell translocation gene 2,rhombotin 2,rhombotin-like 1

Related Disease

[Genetic Predisposition to Disease](#) [Hematologic Diseases](#) [Hodgkin Disease](#)
[Lung Neoplasms](#) [Lymphoma, B-Cell](#) [Lymphoma, Follicular](#)
[Lymphoma, Large B-Cell, Diffuse](#) [Lymphoproliferative Disorders](#) [Occupational Diseases](#)
[Pulmonary Disease, Chronic Obstructive](#) [Urinary Bladder Neoplasms](#)
[Waldenstrom Macroglobulinemia](#) [Werner syndrome](#)

[服務條款](#) | [隱私權政策](#) | [著作及商標](#) | [網站地圖](#)

©2017 亞諾法生技股份有限公司 Abnova Corporation. 版權所有.