



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

CDKN1B FISH Probe

Catalog # : FA0305

規格 : [200 uL]

List All

Specification

Product Description:	Made to order FISH probes for identification of gene amplification using Fluorescent In Situ Hybridization Technique. (Technology)
Supplied Product:	DAPI Counterstain (1500 ng/mL) 250 uL
Storage Instruction:	Store at 4°C in the dark.
Origin:	Human
Source:	Genomic DNA
Notice:	We strongly recommend the customer to use FFPE FISH PreTreatment Kit 1 (Catalog #: KA2375 or KA2691) for the pretreatment of Formalin-Fixed Paraffin-Embedded (FFPE) tissue sections.
Regulation Status:	For research use only (RUO)

Application Image

Fluorescent In Situ Hybridization (Cell)

Applications

Fluorescent In Situ Hybridization (Cell)

 [Protocol Download](#)

Gene Information

Entrez GeneID: [1027](#)

Gene Name: CDKN1B

Gene Alias: CDKN4,KIP1,MEN1B,MEN4,P27KIP1

Gene Description: cyclin-dependent kinase inhibitor 1B (p27, Kip1)

Omim ID: [600778](#), [610755](#)

Gene Ontology: [Hyperlink](#)

Gene Summary: This gene encodes a cyclin-dependent kinase inhibitor, which shares a limited similarity with CDK inhibitor CDKN1A/p21. The encoded protein binds to and prevents the activation of cyclin E-CDK2 or cyclin D-CDK4 complexes, and thus controls the cell cycle progression at G1. The degradation of this protein, which is triggered by its CDK dependent phosphorylation and subsequent ubiquitination by SCF complexes, is required for the cellular transition from quiescence to the proliferative state. [provided by RefSeq]

Other Designations: cyclin-dependent kinase inhibitor 1B

Gene Pathway

[Cell cycle](#) [Chronic myeloid leukemia](#) [ErbB signaling pathway](#) [Pathways in cancer](#)
[Prostate cancer](#) [Small cell lung cancer](#)

Related Disease

[Acromegaly](#) [Adenocarcinoma](#) [Adenocarcinoma, Follicular](#) [Alzheimer Disease](#)
[Alzheimer disease](#) [Ataxia telangiectasia](#) [Atherosclerosis](#) [Breast cancer](#) [Breast Neoplasms](#)
[Carcinoma](#) [Carcinoma, Ductal, Breast](#) [Carcinoma, Hepatocellular](#)
[Carcinoma, Papillary, Follicular](#) [Carcinoma, Squamous Cell](#) [Cardiovascular Diseases](#)
[Chromosome Aberrations](#) [Cognition](#) [Colon cancer](#) [Colorectal Neoplasms](#)

[... see more](#)

[服務條款](#) | [隱私權政策](#) | [著作及商標](#) | [網站地圖](#)

©2017 亞諾法生技股份有限公司 Abnova Corporation. 版權所有.