



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

STAT5A FISH Probe

Catalog # : FA0389

規格 : [200 uL]

List All

Specification

Product Description:	Made to order FISH probes for identification of gene amplification using Fluorescent In Situ Hybridization Technique. (Technology)
Supplied Product:	DAPI Counterstain (1500 ng/mL) 250 uL
Storage Instruction:	Store at 4°C in the dark.
Origin:	Human
Source:	Genomic DNA
Notice:	We strongly recommend the customer to use FFPE FISH PreTreatment Kit 1 (Catalog #: KA2375 or KA2691) for the pretreatment of Formalin-Fixed Paraffin-Embedded (FFPE) tissue sections.
Regulation Status:	For research use only (RUO)

Application Image

Fluorescent In Situ Hybridization (Cell)

Applications

Fluorescent In Situ Hybridization (Cell)

 [Protocol Download](#)

Gene Information

Entrez GeneID: [6776](#)

Gene Name: STAT5A

Gene Alias: MGF,STAT5

Gene Description: signal transducer and activator of transcription 5A

Omim ID: [601511](#)


Gene Ontology: [Hyperlink](#)

Gene Summary: The protein encoded by this gene is a member of the STAT family of transcription factors. In response to cytokines and growth factors, STAT family members are phosphorylated by the receptor associated kinases, and then form homo- or heterodimers that translocate to the cell nucleus where they act as transcription activators. This protein is activated by, and mediates the responses of many cell ligands, such as IL2, IL3, IL7 GM-CSF, erythropoietin, thrombopoietin, and different growth hormones. Activation of this protein in myeloma and lymphoma associated with a TEL/JAK2 gene fusion is independent of cell stimulus and has been shown to be essential for the tumorigenesis. The mouse counterpart of this gene is found to induce the expression of BCL2L1/BCL-X(L), which suggests the antiapoptotic function of this

gene in cells. [provided by RefSeq]

Other -
Designations:

Interactome



Gene Pathway

[Acute myeloid leukemia](#) [Chronic myeloid leukemia](#) [ErbB signaling pathway](#)
[Jak-STAT signaling pathway](#) [Pathways in cancer](#)

Related Disease

[Asthma](#) [Asthma](#) [Birth Weight](#) [Breast cancer](#) [Breast Neoplasms](#) [Bronchiolitis, Viral](#)
[Carcinoma, Hepatocellular](#) [Disease Progression](#) [Genetic Predisposition to Disease](#)
[Glioblastoma](#) [Glioma](#) [Hepatitis C](#) [Infant, Premature, Diseases](#) [Leukemia](#)
[Leukemia, Myeloid, Acute](#) [Liver Neoplasms](#) [Lupus Erythematosus, Systemic](#)
[Meningeal Neoplasms](#) [Meningioma](#)

... see more

[服務條款](#) | [隱私權政策](#) | [著作及商標](#) | [網站地圖](#)

©2017 亞諾法生技股份有限公司 Abnova Corporation. 版權所有.