



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

TNFRSF1B(Texas Red)/CEN1p(FITC) FISH Probe

Catalog # : FA0481

規格 : [200 uL]

[List All](#)

Specification

Product Description:	Made to order FISH probes for identification of gene amplification using Fluorescent In Situ Hybridization Technique. (Technology)
Supplied Product:	DAPI Counterstain (1500 ng/mL) 250 uL
Storage Instruction:	Store at 4°C in the dark.
Origin:	Human
Source:	Genomic DNA
Notice:	We strongly recommend the customer to use FFPE FISH PreTreatment Kit 1 (Catalog #: KA2375 or KA2691) for the pretreatment of Formalin-Fixed Paraffin-Embedded (FFPE) tissue sections.
Regulation Status:	For research use only (RUO)

Application Image

Fluorescent In Situ Hybridization (Cell)

Applications

Fluorescent In Situ Hybridization (Cell)

 [Protocol Download](#)

Gene Information

Entrez GeneID: [7133](#)

Gene Name: TNFRSF1B

Gene Alias: CD120b, TBPII, TNF-R-II, TNF-R75, TNFBR, TNFR1B, TNFR2, TNFR80, p75, p75TNFR

Gene Description: tumor necrosis factor receptor superfamily, member 1B

Omim ID: [191191](#)

Gene Ontology: [Hyperlink](#)

Gene Summary: The protein encoded by this gene is a member of the TNF-receptor superfamily. This protein and TNF-receptor 1 form a heterocomplex that mediates the recruitment of two anti-apoptotic proteins, c-IAP1 and c-IAP2, which possess E3 ubiquitin ligase activity. The function of IAPs in TNF-receptor signalling is unknown, however, c-IAP1 is thought to potentiate TNF-induced apoptosis by the ubiquitination and degradation of TNF-receptor-associated factor 2, which mediates anti-apoptotic signals. Knockout studies in mice also suggest a role of this protein in protecting neurons from apoptosis by stimulating antioxidative pathways. [provided by RefSeq]

Other OTTHUMP00000001784,p75 TNF receptor,soluble TNFR1B variant
Designations: 1,tumor necrosis factor beta receptor,tumor necrosis factor binding protein 2,tumor necrosis factor receptor 2

Gene Pathway

[Adipocytokine signaling pathway](#) [Amyotrophic lateral sclerosis \(ALS\)](#)
[Cytokine-cytokine receptor interaction](#)

Related Disease

[Acute Disease](#) [Adenocarcinoma](#) [Adrenal Hyperplasia, Congenital](#)
[Adrenal hyperplasia, congenital](#) [Aggressive Periodontitis](#) [Albuminuria](#) [Alveolar Bone Loss](#)
[Alzheimer Disease](#) [Alzheimer disease](#) [Anemia](#) [Anemia, Aplastic Anemia, Iron-Deficiency](#)
[Arthritis, Juvenile Rheumatoid Arthritis, Psoriatic Arthritis, Rheumatoid](#) [Aspergillosis](#) [Asthma](#)
[Asthma](#) [Atherosclerosis](#)

... see more

[服務條款](#) | [隱私權政策](#) | [著作及商標](#) | [網站地圖](#)

©2017 亞諾法生技股份有限公司 Abnova Corporation. 版權所有.