



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## EPHA8(Texas Red)/CEN1p(FITC) FISH Probe

Catalog # : FA0482

規格 : [ 200 uL ]

List All

### Specification

<b>Product Description:</b>	Made to order FISH probes for identification of gene amplification using Fluorescent In Situ Hybridization Technique. ( <a href="#">Technology</a> )
<b>Supplied Product:</b>	DAPI Counterstain (1500 ng/mL ) 250 uL
<b>Storage Instruction:</b>	Store at 4°C in the dark.
<b>Origin:</b>	Human
<b>Source:</b>	Genomic DNA
<b>Notice:</b>	We <b>strongly recommend</b> the customer to use FFPE FISH PreTreatment Kit 1 (Catalog #: <a href="#">KA2375</a> or <a href="#">KA2691</a> ) for the pretreatment of Formalin-Fixed Paraffin-Embedded (FFPE) tissue sections.
<b>Regulation Status:</b>	For research use only (RUO)

### Application Image

Fluorescent In Situ Hybridization (Cell)

### Applications

Fluorescent In Situ Hybridization (Cell)

 [Protocol Download](#)

### Gene Information

Entrez GeneID: [2046](#)

Gene Name: EPHA8

Gene Alias: EEK,HEK3,KIAA1459

Gene Description: EPH receptor A8

Omim ID: [176945](#)

Gene Ontology: [Hyperlink](#)

**Gene Summary:** This gene encodes a member of the ephrin receptor subfamily of the protein-tyrosine kinase family. EPH and EPH-related receptors have been implicated in mediating developmental events, particularly in the nervous system. Receptors in the EPH subfamily typically have a single kinase domain and an extracellular region containing a Cys-rich domain and 2 fibronectin type III repeats. The ephrin receptors are divided into 2 groups based on the similarity of their extracellular domain sequences and their affinities for binding ephrin-A and ephrin-B ligands. The protein encoded by this gene functions as a receptor for ephrin A2, A3 and A5 and plays a role in short-range contact-mediated axonal guidance during development of the mammalian nervous system. [provided by RefSeq]

**Other** EPH- and ELK-related tyrosine  
**Designations:** kinase,OTTHUMP0000002934,OTTHUMP0000002935,ephrin type-A  
receptor 8,hydroxyaryl-protein kinase,protein-tyrosine kinase,tyrosine-  
protein kinase receptor EEK,tyrosylprotein kinase

---

### Gene Pathway

---

[Axon guidance](#)

### Related Disease

---

[Celiac Disease](#) [Genetic Predisposition to Disease](#)

---

[服務條款](#) | [隱私權政策](#) | [著作及商標](#) | [網站地圖](#)

©2017 亞諾法生技股份有限公司 Abnova Corporation. 版權所有.