



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com



[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

FGR(Texas Red)/CEN1p(FITC) FISH Probe

Catalog # : FA0485

規格 : [200 uL]

List All

Specification	Application Image
Product Description: Made to order FISH probes for identification of gene amplification using Fluorescent In Situ Hybridization Technique. (Technology)	
Supplied Product: DAPI Counterstain (1500 ng/mL) 250 uL	
Storage Instruction: Store at 4°C in the dark.	
Origin: Human	
Source: Genomic DNA	
Notice: We strongly recommend the customer to use FFPE FISH PreTreatment Kit 1 (Catalog #: KA2375 or KA2691) for the pretreatment of Formalin-Fixed Paraffin-Embedded (FFPE) tissue sections.	
Regulation Status: For research use only (RUO)	
<h3 style="margin: 0;">Applications</h3>	
Fluorescent In Situ Hybridization (Cell)	
 Protocol Download	
<h3 style="margin: 0;">Gene Information</h3>	
Entrez GeneID: 2268	
Gene Name: FGR	
Gene Alias: FLJ43153,MGC75096, SRC2,c-fgr,c-src2,p55c-fgr,p58c-fgr	
Gene Description: Gardner-Rasheed feline sarcoma viral (v-fgr) oncogene homolog	
Omim ID: 164940	
Gene Ontology: Hyperlink	
Gene Summary: This gene is a member of the Src family of protein tyrosine kinases (PTKs). The encoded protein contains N-terminal sites for myristylation and palmitoylation, a PTK domain, and SH2 and SH3 domains which are involved in mediating protein-protein interactions with phosphotyrosine-containing and proline-rich motifs, respectively. The protein localizes to plasma membrane ruffles, and functions as a negative regulator of cell migration and adhesion triggered by the beta-2 integrin signal transduction pathway. Infection with Epstein-Barr virus results in the overexpression of this gene. Multiple alternatively spliced variants, encoding the same protein, have been identified. [provided by RefSeq]	
Other OTTHUMP00000003601,OTTHUMP00000003602,OTTHUMP000000003	

Designations: 603,c-fgr protooncogene,c-src-2 proto-oncogene,p55-c-fgr protein,proto-oncogene tyrosine-protein kinase FGR

Gene Pathway

Chemokine signaling pathway

Related Disease

HIV Infections

[服務條款](#) | [隱私權政策](#) | [著作及商標](#) | [網站地圖](#)

©2017 亞諾法生技股份有限公司 Abnova Corporation. 版權所有.