



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Eph A7(Texas Red)/CEN6p(FITC) FISH Probe

Catalog # : FA0530

規格 : [200 uL]

List All

Specification

Product Description:	Made to order FISH probes for identification of gene amplification using Fluorescent In Situ Hybridization Technique. (Technology)
Supplied Product:	DAPI Counterstain (1500 ng/mL) 250 uL
Storage Instruction:	Store at 4°C in the dark.
Origin:	Human
Source:	Genomic DNA
Notice:	We strongly recommend the customer to use FFPE FISH PreTreatment Kit 1 (Catalog #: KA2375 or KA2691) for the pretreatment of Formalin-Fixed Paraffin-Embedded (FFPE) tissue sections.
Regulation Status:	For research use only (RUO)

Application Image

Fluorescent In Situ Hybridization (Cell)

Applications

Fluorescent In Situ Hybridization (Cell)

 [Protocol Download](#)

Gene Information

Entrez GeneID: [2045](#)

Gene Name: EPHA7

Gene Alias: EHK3,HEK11

Gene Description: EPH receptor A7

Omim ID: [602190](#)

Gene Ontology: [Hyperlink](#)

Gene Summary: This gene belongs to the ephrin receptor subfamily of the protein-tyrosine kinase family. EPH and EPH-related receptors have been implicated in mediating developmental events, particularly in the nervous system. Receptors in the EPH subfamily typically have a single kinase domain and an extracellular region containing a Cys-rich domain and 2 fibronectin type III repeats. The ephrin receptors are divided into 2 groups based on the similarity of their extracellular domain sequences and their affinities for binding ephrin-A and ephrin-B ligands. [provided by RefSeq]

Other Designations: Eph homology kinase-3,OTTHUMP00000016875,OTTHUMP00000040586,ephrin receptor

EphA7, ephrin type-A receptor 7, receptor protein-tyrosine kinase
HEK11, tyrosine-protein kinase receptor EHK-3

Gene Pathway

[Axon guidance](#)

Related Disease

[Tobacco Use Disorder](#)

[服務條款](#) | [隱私權政策](#) | [著作及商標](#) | [網站地圖](#)

©2017 亞諾法生技股份有限公司 Abnova Corporation. 版權所有.