



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)

ARMET(Texas Red)/CEN3q(FITC) FISH Probe

Catalog # : FA0620

規格 : [200 uL]

List All

Specification

Product Description:	Made to order FISH probes for identification of gene amplification using Fluorescent In Situ Hybridization Technique. (Technology)
Supplied Product:	DAPI Counterstain (1500 ng/mL) 250 uL
Storage Instruction:	Store at 4°C in the dark.
Origin:	Human
Source:	Genomic DNA
Notice:	We strongly recommend the customer to use FFPE FISH PreTreatment Kit 1 (Catalog #: KA2375 or KA2691) for the pretreatment of Formalin-Fixed Paraffin-Embedded (FFPE) tissue sections.
Regulation Status:	For research use only (RUO)

Application Image

Fluorescent In Situ Hybridization (Cell)

Applications

Fluorescent In Situ Hybridization (Cell)

 [Protocol Download](#)

Gene Information

Entrez GeneID: [7873](#)

Gene Name: ARMET

Gene Alias: ARP,MANF,MGC142148,MGC142150

Gene Description: arginine-rich, mutated in early stage tumors

Omim ID: [260350](#), [601916](#)

Gene Ontology: [Hyperlink](#)

Gene Summary: The protein encoded by this gene is localized in the endoplasmic reticulum (ER) and golgi, and is also secreted. Reducing expression of this gene increases susceptibility to ER stress-induced death and promotes cell proliferation. The protein was initially thought to be longer at the N-terminus and to contain an arginine-rich region but transcribed evidence indicates a smaller open reading frame that does not encode the arginine tract. The presence of a specific mutation changing the previously numbered codon 50 from ATG to AGG, or deletion of that codon, has been reported in a variety of solid tumors. With the protein size correction, this codon is now identified as the initiation codon. [provided by RefSeq]

Other arginine-rich protein
Designations:

[服務條款](#) | [隱私權政策](#) | [著作及商標](#) | [網站地圖](#)

©2017 亞諾法生技股份有限公司 Abnova Corporation. 版權所有.