



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)

## Datasheet

### H2AFX (Human) Recombinant Protein (P01)

**Catalog Number:** H00003014-P01

**Regulation Status:** For research use only (RUO)

**Product Description:** Human H2AFX full-length ORF (AAH04915, 1 a.a. - 143 a.a.) recombinant protein with GST-tag at N-terminal.

**Sequence:**

MSGRGKTGGKARAKAKSRSSRAGLQFPVGRVHLLR  
KGHYAERVGAGAPVYLAADVLEYLTAEILELAGNAARDN  
KKTRIIIPRHLQLAIRNDEELNLLGGVTIAQGGVLPNIQA  
VLLPKKTSATVGPAPSGGKKATQASQEY

**Host:** Wheat Germ (in vitro)

**Theoretical MW (kDa):** 41.47

**Applications:** AP, Array, ELISA, WB-Re

(See our web site product page for detailed applications information)

**Protocols:** See our web site at

<http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

**Preparation Method:** [in vitro wheat germ expression system](#)

**Purification:** Glutathione Sepharose 4 Fast Flow

**Storage Buffer:** 50 mM Tris-HCl, 10 mM reduced Glutathione, pH=8.0 in the elution buffer.

**Storage Instruction:** Store at -80°C. Aliquot to avoid repeated freezing and thawing.

**Entrez GeneID:** 3014

**Gene Symbol:** H2AFX

**Gene Alias:** H2A.X, H2A/X, H2AX

**Gene Summary:** Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, and H4)

form an octamer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. The linker histone, H1, interacts with linker DNA between nucleosomes and functions in the compaction of chromatin into higher order structures. This gene encodes a member of the histone H2A family, and generates two transcripts through the use of the conserved stem-loop termination motif, and the polyA addition motif. [provided by RefSeq]