



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)

Datasheet

NDUFB7 (Human) Recombinant Protein (P01)

Catalog Number: H00004713-P01

Regulation Status: For research use only (RUO)

Product Description: Human NDUFB7 full-length ORF (NP_004137.2, 1 a.a. - 137 a.a.) recombinant protein with GST-tag at N-terminal.

Sequence:

MG AHLVRRYLGDASVEPDPLQMPTFPPDYGFPERKE
RE MVATQQEMMDAQLRLQLRDYCAHHLIRLLKCKRDS
FPNFLACKQERHDWDYCEHRDYVMRMKEFERERLL
QRKKRREKKAELAKGQGPGEVDPKVAL

Host: Wheat Germ (in vitro)

Theoretical MW (kDa): 42.8

Applications: AP, Array, ELISA, WB-Re

(See our web site product page for detailed applications information)

Protocols: See our web site at

<http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Preparation Method: [in vitro wheat germ expression system](#)

Purification: Glutathione Sepharose 4 Fast Flow

Storage Buffer: 50 mM Tris-HCl, 10 mM reduced Glutathione, pH=8.0 in the elution buffer.

Storage Instruction: Store at -80°C. Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 4713

Gene Symbol: NDUFB7

Gene Alias: B18, CI-B18, MGC2480

Gene Summary: The protein encoded by this gene is a subunit of the multisubunit NADH:ubiquinone oxidoreductase (complex I). Mammalian complex I is composed of 45 different subunits. It is located at the

mitochondrial inner membrane. This protein has NADH dehydrogenase activity and oxidoreductase activity. It transfers electrons from NADH to the respiratory chain. The immediate electron acceptor for the enzyme is believed to be ubiquinone. [provided by RefSeq]