



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

POMC (Human) Recombinant Protein (P02)

Catalog Number: H00005443-P02

Regulation Status: For research use only (RUO)

Product Description: Human POMC full-length ORF (NP_000930.1, 1 a.a. - 267 a.a.) recombinant protein with GST-tag at N-terminal.

Sequence:

MPRSCCSRSGALLLALLLQASMEVRGWCLESSQCQD
LTESNLLECIRACKPDLAETPMFPGNGDEQPLTENP
RKYVMGHFRWDRFRNRNSSSSGSSGAGQKREDVSA
GEDCGPLPEGGPEPRSDGAKPGPREGKRSYSMEHF
RWGKPVGKKRRPVKVPNGAEDESAAFPLEFKREL
TGQRLREGDGPDPADGAGAQAADLEHSLLVAAEKK
DEGYPYRMEHFRWGSPPKDKRYGGFMTSEKSQTPLVT
LFKNAIIKNAYKKGE

Host: Wheat Germ (in vitro)

Theoretical MW (kDa): 55.8

Applications: AP, Array, ELISA, WB-Re
(See our web site product page for detailed applications information)

Protocols: See our web site at
<http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Preparation Method: [in vitro wheat germ expression system](#)

Purification: Glutathione Sepharose 4 Fast Flow

Storage Buffer: 50 mM Tris-HCl, 10 mM reduced Glutathione, pH=8.0 in the elution buffer.

Storage Instruction: Store at -80°C. Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 5443

Gene Symbol: POMC

Gene Alias: ACTH, CLIP, LPH, MSH, NPP, POC

Gene Summary: This gene encodes a polypeptide hormone precursor that undergoes extensive, tissue-specific, post-translational processing via cleavage by subtilisin-like enzymes known as prohormone convertases. There are eight potential cleavage sites within the polypeptide precursor and, depending on tissue type and the available convertases, processing may yield as many as ten biologically active peptides involved in diverse cellular functions. The encoded protein is synthesized mainly in corticotroph cells of the anterior pituitary where four cleavage sites are used; adrenocorticotrophin, essential for normal steroidogenesis and the maintenance of normal adrenal weight, and lipotropin beta are the major end products. In other tissues, including the hypothalamus, placenta, and epithelium, all cleavage sites may be used, giving rise to peptides with roles in pain and energy homeostasis, melanocyte stimulation, and immune modulation. These include several distinct melanotropins, lipotropins, and endorphins that are contained within the adrenocorticotrophin and beta-lipotropin peptides. Mutations in this gene have been associated with early onset obesity, adrenal insufficiency, and red hair pigmentation. Alternatively spliced transcript variants encoding the same protein have been described. [provided by RefSeq]