



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

SDCBP monoclonal antibody (MA01), clone 2C12 (APC)

Catalog Number: H00006386-MA01

Regulation Status: For research use only (RUO)

Product Description: APC conjugated mouse monoclonal antibody raised against a partial recombinant SDCBP.

Clone Name: 2C12

Immunogen: SDCBP (NP_005616, 1 a.a. ~ 100 a.a)
partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Sequence:

MSLYPSLEDLKVDKVIQAQTAFSANPANPAILSEASAPI
PHDGNLYPRLYPELSQYMGLSLNEEEIRANVAVVSGA
PLQGQLVARPSSINYMVAPVTGND

Host: Mouse

Protocols: See our web site at
<http://www.abnova.com/support/protocols.asp> or product
page for detailed protocols

Conjugation: APC

Isotype: IgG1 Kappa

Storage Buffer: In 1x PBS, pH 7.4

Storage Instruction: Store at -20°C or lower. Aliquot to
avoid repeated freezing and thawing.

Entrez GeneID: 6386

Gene Symbol: SDCBP

Gene Alias: MDA-9, ST1, SYCL, TACIP18

Gene Summary: The protein encoded by this gene was initially identified as a molecule linking syndecan-mediated signaling to the cytoskeleton. The syntenin protein contains tandemly repeated PDZ domains that bind the cytoplasmic, C-terminal domains of a variety of transmembrane proteins. This protein may also affect cytoskeletal-membrane organization, cell adhesion,

protein trafficking, and the activation of transcription factors. The protein is primarily localized to membrane-associated adherens junctions and focal adhesions but is also found at the endoplasmic reticulum and nucleus. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq]