



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

TNFRSF10C (Human) Recombinant Protein (P01)

Catalog Number: H00008794-P01

Regulation Status: For research use only (RUO)

Product Description: Human TNFRSF10C full-length ORF (NP_003832.2, 1 a.a. - 259 a.a.) recombinant protein with GST-tag at N-terminal.

Sequence:

```
MARIPKTLKFVVVIVAVLLPVLAYSATTARQEEVPQQTV  
APQQQRHSFKGEECPAGSHRSEHTGACNPCTEGVDY  
TNASNNEPSCFPCTVCKSDQKHKSCTMTRDTCQC  
KEGTFRNENSPMCRKCSRCPGSEVQVSNCTSWDDI  
QCVEEFGANATVETPAAEETMNTSPGTPAPAAEETMN  
TSPGTPAPAAEETMTTSPGTPAPAAEETMTTSPGTPA  
PAAEETMTTSPGTPASSHYLSCTIVGIIVLIVLLIVFV
```

Host: Wheat Germ (in vitro)

Theoretical MW (kDa): 53.8

Applications: AP, Array, ELISA, WB-Re
(See our web site product page for detailed applications information)

Protocols: See our web site at
<http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Preparation Method: [in vitro wheat germ expression system](#)

Purification: Glutathione Sepharose 4 Fast Flow

Storage Buffer: 50 mM Tris-HCl, 10 mM reduced Glutathione, pH=8.0 in the elution buffer.

Storage Instruction: Store at -80°C. Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 8794

Gene Symbol: TNFRSF10C

Gene Alias: CD263, DCR1, LIT, MGC149501, MGC149502, TRAILR3, TRID

Gene Summary: The protein encoded by this gene is a member of the TNF-receptor superfamily. This receptor contains an extracellular TRAIL-binding domain and a transmembrane domain, but no cytoplasmic death domain. This receptor is not capable of inducing apoptosis, and is thought to function as an antagonistic receptor that protects cells from TRAIL-induced apoptosis. This gene was found to be a p53-regulated DNA damage-inducible gene. The expression of this gene was detected in many normal tissues but not in most cancer cell lines, which may explain the specific sensitivity of cancer cells to the apoptosis-inducing activity of TRAIL. [provided by RefSeq]