



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

BET1 (Human) Recombinant Protein (P01)

Catalog Number: H00010282-P01

Regulation Status: For research use only (RUO)

Product Description: Human BET1 full-length ORF (AAH00899, 1 a.a. - 118 a.a.) recombinant protein with GST-tag at N-terminal.

Sequence:

MRRAGLGEGVPPGNYGNYGYANSGYSACEEENERLT
ESLRKVTAIKSLISIEIGHEVKTQNKLLAEMDSQFDSTT
GFLGKTMGKILSRGSQTKLLCYMMLFSLFVFFIYWII
KLR

Host: Wheat Germ (in vitro)

Theoretical MW (kDa): 38.72

Interspecies Antigen Sequence: Mouse (92); Rat (93)

Applications: AP, Array, ELISA, WB-Re

(See our web site product page for detailed applications information)

Protocols: See our web site at

<http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Preparation Method: [in vitro wheat germ expression system](#)

Purification: Glutathione Sepharose 4 Fast Flow

Storage Buffer: 50 mM Tris-HCl, 10 mM reduced Glutathione, pH=8.0 in the elution buffer.

Storage Instruction: Store at -80°C. Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 10282

Gene Symbol: BET1

Gene Alias: DKFZp781C0425, HBET1

Gene Summary: This gene encodes a golgi-associated membrane protein that participates in vesicular transport

from the endoplasmic reticulum (ER) to the Golgi complex. The encoded protein functions as a soluble N-ethylmaleimide-sensitive factor attachment protein receptor and may be involved in the docking of ER-derived vesicles with the cis-Golgi membrane. Alternatively spliced transcript variants encoding different isoforms have been described but their full-length nature has not been determined. [provided by RefSeq]