



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

CCL27 (Human) Recombinant Protein (P01)

Catalog Number: H00010850-P01

Regulation Status: For research use only (RUO)

Product Description: Human CCL27 full-length ORF (NP_006655, 1 a.a. - 112 a.a.) recombinant protein with GST-tag at N-terminal.

Sequence:

MKGPPTFCSLLLSLLLSPDPTAAFLPPSTACCTQLY
RKPLSDKLLRKVIQVELQEADGDCHLQAFVLHLAQRSI
CIHPQNPSLSQWFEHQERKLHGTLPKLNFGMLRKMKG

Host: Wheat Germ (in vitro)

Theoretical MW (kDa): 37.95

Interspecies Antigen Sequence: Mouse (61)

Applications: AP, Array, ELISA, WB-Re
(See our web site product page for detailed applications information)

Protocols: See our web site at
<http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Preparation Method: [in vitro wheat germ expression system](#)

Purification: Glutathione Sepharose 4 Fast Flow

Storage Buffer: 50 mM Tris-HCl, 10 mM reduced Glutathione, pH=8.0 in the elution buffer.

Storage Instruction: Store at -80°C. Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 10850

Gene Symbol: CCL27

Gene Alias: ALP, CTACK, CTAK, ESKINE, ILC, PESKY, SCYA27

Gene Summary: This gene is one of several CC cytokine genes clustered on the p-arm of chromosome

9. Cytokines are a family of secreted proteins involved in immunoregulatory and inflammatory processes. The CC cytokines are proteins characterized by two adjacent cysteines. The protein encoded by this gene is chemotactic for skin-associated memory T lymphocytes. This cytokine may also play a role in mediating homing of lymphocytes to cutaneous sites. It specifically binds to chemokine receptor 10 (CCR10). Studies of a similar murine protein indicate that these protein-receptor interactions have a pivotal role in T cell-mediated skin inflammation. [provided by RefSeq]