



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

WBP2 (Human) Recombinant Protein (P01)

Catalog Number: H00023558-P01

Regulation Status: For research use only (RUO)

Product Description: Human WBP2 full-length ORF (AAH10616, 1 a.a. - 261 a.a.) recombinant protein with GST-tag at N-terminal.

Sequence:

```
MALNKNHSEGGGVIVNNTESILMSYDHVELTFNDMKN  
VPEAFKGTKKGTVYLTPYRVIFLSKGDAMQSFMMPF  
YLMKDCEIKQPVFGANYIKGTVKAEAGGGWEGSASYK  
LTFTAGGAIEFGQRMLQVASQASRGEVPSGAYGYSY  
MPSGAYVYPPPVANGMYPCPPGYPPPPPEFYPGP  
PMDGAMGYVQPPPPYPGMEPPVSGPDVPSTPA  
AEAKAAEAAASAYNPGNPHNVYMPTSQPPPPYP  
PEDKKTQ
```

Host: Wheat Germ (in vitro)

Theoretical MW (kDa): 54.45

Interspecies Antigen Sequence: Mouse (97); Rat (96)

Applications: AP, Array, ELISA, WB-Re

(See our web site product page for detailed applications information)

Protocols: See our web site at

<http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Preparation Method: [in vitro wheat germ expression system](#)

Purification: Glutathione Sepharose 4 Fast Flow

Storage Buffer: 50 mM Tris-HCl, 10 mM reduced Glutathione, pH=8.0 in the elution buffer.

Storage Instruction: Store at -80°C. Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 23558

Gene Symbol: WBP2

Gene Alias: MGC18269, WBP-2

Gene Summary: The globular WW domain is composed of 38 to 40 semiconserved amino acids shared by proteins of diverse functions including structural, regulatory, and signaling proteins. The domain is involved in mediating protein-protein interactions through the binding of polyproline ligands. This gene encodes a WW domain binding protein, which binds to the WW domain of Yes kinase-associated protein by its PY motifs. The function of this protein has not been determined. [provided by RefSeq]