



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

IL17D (Human) Recombinant Protein (P01)

Catalog Number: H00053342-P01

Regulation Status: For research use only (RUO)

Product Description: Human IL17D full-length ORF (NP_612141.1, 1 a.a. - 202 a.a.) recombinant protein with GST-tag at N-terminal.

Sequence:

MLVAGFLLALPPSWAAGAPRAGRPPARPRGCADRPE
ELLEQLYGRLAAGVLSAFHHTLQLGPREQARNASCPA
GGRPADRRFRPPTNLRVSPWAYRISYDPARYPRYLP
EAYCLCRGCLTGLFGCEEDVFRSAPVYMPTVLLR RTP
ACAGRSVYTEAYVTIPVGCTCVPEPEKDADSINSSID
KQGAKLLLGPNDAPAGP

Host: Wheat Germ (in vitro)

Theoretical MW (kDa): 48.3

Interspecies Antigen Sequence: Mouse (85)

Applications: AP, Array, ELISA, WB-Re
(See our web site product page for detailed applications information)

Protocols: See our web site at
<http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Preparation Method: [in vitro wheat germ expression system](#)

Purification: Glutathione Sepharose 4 Fast Flow

Storage Buffer: 50 mM Tris-HCl, 10 mM reduced Glutathione, pH=8.0 in the elution buffer.

Storage Instruction: Store at -80°C. Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 53342

Gene Symbol: IL17D

Gene Alias: FLJ30846, IL-17D, IL-22, IL-27, IL27

Gene Summary: The protein encoded by this gene is a cytokine that shares the sequence similarity with IL17. The treatment of endothelial cells with this cytokine has been shown to stimulate the production of other cytokines including IL6, IL8 and CSF2/ GM-CSF. The increased expression of IL8 induced by this cytokine was found to be NF-kappa B-dependent. [provided by RefSeq]