



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

WDR45L (Human) Recombinant Protein (P01)

Catalog Number: H00056270-P01

Regulation Status: For research use only (RUO)

Product Description: Human WDR45L full-length ORF (AAH07838.1, 1 a.a. - 286 a.a.) recombinant protein with GST-tag at N-terminal.

Sequence:

MLFRCNYLALVGGGKKPKYPPNKVMIWDDLKKTVEI
EFSTEVKAVKLRRDRIVVLDSEMIKVFTHNPHQLHV
FETCYNPKGLCVLCPNSNLLAFPGTHTGHVQLVDL
ASTEKPPVDIPAHEGVLSCIALNLQGTRITASEKGTLI
RIFDTSSGHLIQELRRGSQAANIYCINFNQDASLICVSS
DHGTVHIFAAEDPKRNKQSSLASASFLPKYFSSKWSF
SKFQVPSGSPCICAFGTEPNAVIAICADGSYYKFLFNP
KGECIRDVYAQFLEMTDDKL

Host: Wheat Germ (in vitro)

Theoretical MW (kDa): 58.1

Interspecies Antigen Sequence: Mouse (98); Rat (98)

Applications: AP, Array, ELISA, WB-Re
(See our web site product page for detailed applications information)

Protocols: See our web site at
<http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Preparation Method: [in vitro wheat germ expression system](#)

Purification: Glutathione Sepharose 4 Fast Flow

Storage Buffer: 50 mM Tris-HCl, 10 mM reduced Glutathione, pH=8.0 in the elution buffer.

Storage Instruction: Store at -80°C. Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 56270

Gene Symbol: WDR45L

Gene Alias: WIPI-3, WIPI3

Gene Summary: This gene encodes a member of the WIPI or SVP1 family of WD40 repeat-containing proteins. The protein contains seven WD40 repeats that are thought to fold into a beta-propeller structure that mediates protein-protein interactions, and a conserved motif for interaction with phospholipids. The human genome contains several pseudogenes of this gene. [provided by RefSeq]