



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

TAAR8 (Human) Recombinant Protein (P01)

Catalog Number: H00083551-P01

Regulation Status: For research use only (RUO)

Product Description: Human TAAR8 full-length ORF (NP_444508.1, 1 a.a. - 342 a.a.) recombinant protein with GST-tag at N-terminal.

Sequence:

MTSNFSQPVVQLCYEDVNGSCIETPYSPGSRVILYTAF
SFGSLLAVFGNLLVMTSVLHFKQLHSPTNFLIASLACA
DFLVGVTVMLFSMVRTVESCWYFGAKFCTLHSCCDV
AFCYSSVLHLCFICIDRYIVVTDPLVYATKFTVSVSGICI
SVSWILPLTYSGAVFYTGVDNDGLEELVSALNCVGGC
QIIVSQGWVLIDFLLFFIPTLVMILYSKIFLIAKQQAIIET
TSSKVESSSESYKIRVAKRERKAAKTLGVTVLAFVISW
LPYTVDILIDAFMGFLTPAYIYEICCSAYYNSAMNPLIY
ALFYPWFRKAIKLILSGDVLKASSSTISLFLE

Host: Wheat Germ (in vitro)

Theoretical MW (kDa): 64.4

Applications: AP, Array, ELISA, WB-Re
(See our web site product page for detailed applications information)

Protocols: See our web site at
<http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Preparation Method: [in vitro wheat germ expression system](#)

Purification: Glutathione Sepharose 4 Fast Flow

Storage Buffer: 50 mM Tris-HCl, 10 mM reduced Glutathione, pH=8.0 in the elution buffer.

Storage Instruction: Store at -80°C. Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 83551

Gene Symbol: TAAR8

Gene Alias: GPR102, TA5, TAR5, TRAR5

Gene Summary: G protein-coupled receptors (GPCRs, or GPRs) contain 7 transmembrane domains and transduce extracellular signals through heterotrimeric G proteins.[supplied by OMIM]