



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

CLEC12A (Human) Recombinant Protein

Catalog Number: H00160364-G01

Regulation Status: For research use only (RUO)

Product Description: Human CLEC12A full-length ORF (ADR82900.1) recombinant protein without tag.

Sequence:

MSEEVTYADLQFQNSSEMEDIKPEIGKFGKAPPAPSH
VWRPAALFLTLLCLLLIIGLVLASMFHVTLKIEMKMMN
KLQNISEELQRNLSLQMSNMNINNKIRNLSTTLQTIATK
LCRELYSKEQEHEKCKPCPRRWIWHKDSCYFLSDDVQ
TWQESKMACAAQNASLLKINNKNALEFIKSQRSYDY
WLGLSPEEDSTRGMRVDNIINSSAWVIRNAPDLNNMY
CGYINRLVQYYHCTYKQRMICEKMANPVQLGSTYFR
EA

Host: Wheat Germ (in vitro)

Theoretical MW (kDa): 29.2

Applications: AP

(See our web site product page for detailed applications information)

Protocols: See our web site at

<http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Form: Liquid

Preparation Method: [in vitro wheat germ expression system with proprietary liposome technology](#)

Purification: None

Recommend Usage: Heating may cause protein aggregation. Please do not heat this product before electrophoresis.

Storage Buffer: 25 mM Tris-HCl of pH8.0 containing 2% glycerol.

Storage Instruction: Store at -80°C. Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 160364

Gene Symbol: CLEC12A

Gene Alias: CLL-1, CLL1, DCAL-2, MGC70602, MICL

Gene Summary: This gene encodes a member of the C-type lectin/C-type lectin-like domain (CTL/CTLD) superfamily. Members of this family share a common protein fold and have diverse functions, such as cell adhesion, cell-cell signalling, glycoprotein turnover, and roles in inflammation and immune response. The protein encoded by this gene is a negative regulator of granulocyte and monocyte function. Several alternatively spliced transcript variants of this gene have been described, but the full-length nature of some of these variants has not been determined. This gene is closely linked to other CTL/CTLD superfamily members in the natural killer gene complex region on chromosome 12p13. [provided by RefSeq]