



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## Datasheet

### ITGA3 monoclonal antibody, clone VJ1/6 (FITC)

**Catalog Number:** MAB13692

**Regulatory Status:** For research use only (RUO)

**Product Description:** Mouse monoclonal antibody raised against human ITGA3.

**Clone Name:** VJ1/6

**Immunogen:** TNF activated HUVEC cells.

**Host:** Mouse

**Theoretical MW (kDa):** 150

**Reactivity:** Human

**Applications:** Flow Cyt

(See our web site product page for detailed applications information)

**Protocols:** See our web site at

<http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

**Form:** Liquid

**Conjugation:** FITC

**Purification:** Protein A/G purification

**Purity:** >90%

**Isotype:** IgG1

**Recommend Usage:** Flow Cytometry (20  $\mu$ L/ $10^6$  cells)

The optimal working dilution should be determined by the end user.

**Storage Buffer:** In PBS, pH 7.4 (protein stabilizer, 0.09% sodium azide).

**Storage Instruction:** Store in the dark at 4°C.

**Entrez GeneID:** 3675

**Gene Symbol:** ITGA3

**Gene Alias:** CD49C, FLJ34631, FLJ34704, GAP-B3, GAPB3, MSK18, VCA-2, VL3A, VLA3a

**Gene Summary:** The protein encoded by this gene belongs to the family of integrins. Integrins are heterodimeric integral membrane proteins composed of an alpha chain and a beta chain, and function as cell surface adhesion molecules. This gene encodes alpha 3 subunit, which undergoes post-translational cleavage in the extracellular domain to yield disulfide-linked light and heavy chains that join with beta 1 subunit to form an integrin that interacts with many extracellular-matrix proteins. Alternatively spliced transcript variants encoding different isoforms have been identified for this gene. [provided by RefSeq]