



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

CD8A monoclonal antibody, clone 143-44 (PE)

Catalog Number: MAB13760

Regulatory Status: For research use only (RUO)

Product Description: Mouse monoclonal antibody raised against native human CD8A.

Clone Name: 143-44

Immunogen: T-lymphocyte differentiation antigen.

Host: Mouse

Theoretical MW (kDa): 32

Reactivity: Human

Applications: Flow Cyt

(See our web site product page for detailed applications information)

Protocols: See our web site at

<http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Form: Liquid

Conjugation: PE

Purification: Protein A/G purification

Purity: >90%

Isotype: IgG1

Recommend Usage: Flow Cytometry (20 μ L/ 10^6 cells)

The optimal working dilution should be determined by the end user.

Storage Buffer: In PBS, pH 7.4 (protein stabilizer, 0.09% sodium azide).

Storage Instruction: Store in the dark at 4°C. Avoid prolonged exposure to light.

Entrez GeneID: 925

Gene Symbol: CD8A

Gene Alias: CD8, Leu2, MAL, p32

Gene Summary: The CD8 antigen is a cell surface glycoprotein found on most cytotoxic T lymphocytes that mediates efficient cell-cell interactions within the immune system. The CD8 antigen acts as a corepressor with the T-cell receptor on the T lymphocyte to recognize antigens displayed by an antigen presenting cell (APC) in the context of class I MHC molecules. The coreceptor functions as either a homodimer composed of two alpha chains, or as a heterodimer composed of one alpha and one beta chain. Both alpha and beta chains share significant homology to immunoglobulin variable light chains. This gene encodes the CD8 alpha chain isoforms. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq]