



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## Datasheet

### ITGB2 monoclonal antibody, clone GRF1 (FITC)

**Catalog Number:** MAB13823

**Regulatory Status:** For research use only (RUO)

**Product Description:** Mouse monoclonal antibody raised against human ITGB2.

**Clone Name:** GRF1

**Immunogen:** Human acute lymphoblastic leukemia (ALL) T cells.

**Host:** Mouse

**Theoretical MW (kDa):** 98

**Reactivity:** Human

**Applications:** Flow Cyt  
(See our web site product page for detailed applications information)

**Protocols:** See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

**Form:** Liquid

**Conjugation:** FITC

**Purification:** Protein A/G purification

**Purity:** >90%

**Isotype:** IgG1

**Recommend Usage:** Flow Cytometry (20  $\mu$ L/ $10^6$  cells)  
The optimal working dilution should be determined by the end user.

**Storage Buffer:** In PBS, pH 7.4 (protein stabilizer, 0.09% sodium azide).

**Storage Instruction:** Store in the dark at 4°C. Avoid prolonged exposure to light.

**Entrez GeneID:** 3689

**Gene Symbol:** ITGB2

**Gene Alias:** CD18, LAD, LCAMB, LFA-1, MAC-1, MF17, MFI7

**Gene Summary:** The product of this gene belongs to the integrin beta chain family of proteins. Integrins are integral cell-surface proteins composed of an alpha chain and a beta chain. This gene encodes the integrin beta chain beta 2. A given chain may combine with multiple partners resulting in different integrins. For example, beta 2 combines with the alpha L chain to form the integrin LFA-1, and combines with the alpha M chain to form the integrin Mac-1. Integrins are known to participate in cell adhesion as well as cell-surface mediated signalling. Defects in this gene are the cause of leukocyte adhesion deficiency type I (LAD1). Two transcript variants encoding the same protein have been identified for this gene. [provided by RefSeq]