



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

IL2RA monoclonal antibody, clone TP1/6 (APC)

Catalog Number: MAB13854

Regulatory Status: For research use only (RUO)

Product Description: Mouse monoclonal antibody raised against human IL2RA.

Clone Name: TP1/6

Immunogen: PHA activated human T cells.

Host: Mouse

Theoretical MW (kDa): 55

Reactivity: Human

Applications: Flow Cyt

(See our web site product page for detailed applications information)

Protocols: See our web site at

<http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Form: Liquid

Conjugation: APC

Purification: Protein A/G purification

Purity: >90%

Isotype: IgG2b

Recommend Usage: Flow Cytometry (20 μ L/ 10^6 cells)

The optimal working dilution should be determined by the end user.

Storage Buffer: In PBS, pH 7.4 (protein stabilizer, 0.09% sodium azide).

Storage Instruction: Store in the dark at 4°C. Avoid prolonged exposure to light.

Entrez GeneID: 3559

Gene Symbol: IL2RA

Gene Alias: CD25, IDDM10, IL2R, TCGFR

Gene Summary: The interleukin 2 (IL2) receptor alpha (IL2RA) and beta (IL2RB) chains, together with the common gamma chain (IL2RG), constitute the high-affinity IL2 receptor. Homodimeric alpha chains (IL2RA) result in low-affinity receptor, while homodimeric beta (IL2RB) chains produce a medium-affinity receptor. Normally an integral-membrane protein, soluble IL2RA has been isolated and determined to result from extracellular proteolysis. Alternately-spliced IL2RA mRNAs have been isolated, but the significance of each is presently unknown. [provided by RefSeq]