



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

CD79A monoclonal antibody, clone HM57 (FITC)

Catalog Number: MAB15368

Regulatory Status: For research use only (RUO)

Product Description: Mouse monoclonal antibody raised against synthetic peptide of human CD79A.

Clone Name: HM57

Immunogen: A synthetic peptide corresponding to amino acids 202-216 of human CD79A.

Host: Mouse

Theoretical MW (kDa): 47

Reactivity: Human

Applications: Flow Cyt
(See our web site product page for detailed applications information)

Protocols: See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Form: Liquid

Conjugation: FITC

Purification: Affinity purification

Isotype: IgG1

Recommend Usage: Flow Cytometry (20 μ L/ 10^6 cells)
The optimal working dilution should be determined by the end user.

Storage Buffer: In PBS, pH 7.4 (protein stabilizer, 0.09% sodium azide).

Storage Instruction: Store in the dark at 4°C.

Entrez GeneID: 973

Gene Symbol: CD79A

Gene Alias: IGA, MB-1

Gene Summary: The B lymphocyte antigen receptor is a multimeric complex that includes the antigen-specific component, surface immunoglobulin (Ig). Surface Ig non-covalently associates with two other proteins, Ig-alpha and Ig-beta, which are necessary for expression and function of the B-cell antigen receptor. This gene encodes the Ig-alpha protein of the B-cell antigen component. Alternatively spliced transcript variants encoding different isoforms have been described. [provided by RefSeq]