



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

CD8 FITC monoclonal antibody, clone hCD8

Catalog Number: MAB23590

Regulation Status: For research use only (RUO)

Product Description: Mouse FITC monoclonal antibody raised against synthetic peptide of human CD8.

Clone Name: hCD8

Immunogen: Purified human PBL CD8+ T cells

Host: Mouse

Protocols: See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Form: Lyophilized

Purification: Ion exchange column

Isotype: IgG2a

Recommend Usage: weakly cytotoxic. For staining (10 μ L/1,000,000 cells). Titer for blocking T cells should be determined by the investigator.

The optimal working dilution should be determined by the end user.

Storage Buffer: Lyophilized from 1 mg/mL in PBS

Storage Instruction: Store at 4°C on dry atmosphere. After reconstitution with 1 mL of deionized water, store at -20°C or lower.

Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 925

Gene Symbol: CD8A

Gene Alias: CD8, Leu2, MAL, p32

Gene Summary: The CD8 antigen is a cell surface glycoprotein found on most cytotoxic T lymphocytes that mediates efficient cell-cell interactions within the immune system. The CD8 antigen acts as a corepressor with the T-cell receptor on the T lymphocyte to recognize

antigens displayed by an antigen presenting cell (APC) in the context of class I MHC molecules. The coreceptor functions as either a homodimer composed of two alpha chains, or as a heterodimer composed of one alpha and one beta chain. Both alpha and beta chains share significant homology to immunoglobulin variable light chains. This gene encodes the CD8 alpha chain isoforms. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq]