



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

CD44 monoclonal antibody, clone P5C10AT

Catalog Number: MAB23614

Regulation Status: For research use only (RUO)

Product Description: Mouse monoclonal antibody raised against synthetic peptide of human CD44, clone P5C10AT.

Clone Name: P5C10AT

Immunogen: A synthetic peptide corresponding to amino acids 21-145 of human CD44 purified from *Escherichia Coli*.

Host: Mouse

Protocols: See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Form: Liquid

Purification: protein-G affinity chromatography

Isotype: IgG2b heavy chain and ? light chain

Recommend Usage: Western Blot

ELISA

The Western blot analysis is 1:1,000 ~ 2,000.

Recommended starting dilution is 1:1,000.

Storage Buffer: In PBS, pH 7.4 and 0.1% Sodium Azide

Storage Instruction: Store at 4°C for 1 month.

After reconstitution with 1 mL of deionized water, store at -20°C for 12 months.

Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 960

Gene Symbol: CD44

Gene Alias: CDW44, CSPG8, ECMR-III, HCELL, IN, LHR, MC56, MDU2, MDU3, MGC10468, MIC4, MUTCH-I, Pgp1

Gene Summary: The protein encoded by this gene is a

cell-surface glycoprotein involved in cell-cell interactions, cell adhesion and migration. It is a receptor for hyaluronic acid (HA) and can also interact with other ligands, such as osteopontin, collagens, and matrix metalloproteinases (MMPs). This protein participates in a wide variety of cellular functions including lymphocyte activation, recirculation and homing, hematopoiesis, and tumor metastasis. Transcripts for this gene undergo complex alternative splicing that results in many functionally distinct isoforms, however, the full length nature of some of these variants has not been determined. Alternative splicing is the basis for the structural and functional diversity of this protein, and may be related to tumor metastasis. [provided by RefSeq]