

## Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

#### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

#### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com



9F, No. 108, Jhouzih St.,Taipei, Taiwan Tel: + 886-2-8751-1888 Fax: + 886-2-6602-1218 E-mail: sales@abnova.com

### Datasheet

# CDK4 monoclonal antibody, clone P6D10AT

Catalog Number: MAB23652

Regulation Status: For research use only (RUO)

**Product Description:** Mouse monoclonal antibody raised against synthetic peptide of human CDK4, clone P6D10AT.

Clone Name: P6D10AT

**Immunogen:** A synthetic peptide corresponding to amino acids 1-303 of human CDK4 purified from *Escherichia Coli*.

Host: Mouse

**Protocols:** See our web site at http://www.abnova.com/support/protocols.asp or product page for detailed protocols

Form: Liquid

**Purification:** Protein-G affinity chromatography

**Isotype:** Mouse IgG2b heavy chain and kappa light chain

**Storage Buffer:** In PBS, pH 7.4, 10% glycerol and 0.02% Sodium Azide

**Storage Instruction:** Store at 4°C for 1 month. After reconstitution with 1 mL of deionized water, store at -20°C for 12 months. Aliquot to avoid repeated freezing and thawing.

Entrez GenelD: 1019

Gene Symbol: CDK4

Gene Alias: CMM3, MGC14458, PSK-J3

**Gene Summary:** The protein encoded by this gene is a member of the Ser/Thr protein kinase family. This protein is highly similar to the gene products of S. cerevisiae cdc28 and S. pombe cdc2. It is a catalytic subunit of the protein kinase complex that is important for cell cycle G1 phase progression. The activity of this

kinase is restricted to the G1-S phase, which is controlled by the regulatory subunits D-type cyclins and CDK inhibitor p16(INK4a). This kinase was shown to be responsible for the phosphorylation of retinoblastoma gene product (Rb). Mutations in this gene as well as in its related proteins including D-type cyclins, p16(INK4a) and Rb were all found to be associated with tumorigenesis of a variety of cancers. Multiple polyadenylation sites of this gene have been reported. [provided by RefSeq]