



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

LDHA monoclonal antibody, clone 20

Catalog Number: MAB9910

Regulatory Status: For research use only (RUO)

Product Description: Mouse monoclonal antibody raised against full length recombinant LDHA.

Clone Name: 20

Immunogen: Recombinant protein corresponding to full length human LDHA.

Host: Mouse

Reactivity: Human

Applications: ELISA, WB-Ce
(See our web site product page for detailed applications information)

Protocols: See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Specificity: It can expression in HepG2 whole cell lysate.

Form: Liquid

Purification: Affinity purification

Isotype: IgG2b

Recommend Usage: Western blot (1:500)
The optimal working dilution should be determined by the end user.

Storage Buffer: In Citrate-Tris-HCl buffer, pH 7.0
(0.02% Proclin 300)

Storage Instruction: Store at 4°C. For long term storage store at -20°C.
Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 3939

Gene Symbol: LDHA

Gene Alias: LDH-M, LDH1, PIG19

Gene Summary: The protein encoded by this gene catalyzes the conversion of L-lactate and NAD to pyruvate and NADH in the final step of anaerobic glycolysis. The protein is found predominantly in muscle tissue and belongs to the lactate dehydrogenase family. Mutations in this gene have been linked to exertional myoglobinuria. Multiple transcript variants encoding different isoforms have been found for this gene. The human genome contains several non-transcribed pseudogenes of this gene. [provided by RefSeq]