



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



Datasheet

IL15 (Human) Recombinant Protein

Catalog Number: P8346

Regulation Status: For research use only (RUO)

Product Description: Human IL15 (P40933, 48 a.a. - 162 a.a.) partial recombinant protein expressed in *Escherichia coli*.

Sequence:

MNWVNVISDLKKIEDLIQSMHIDATLYTESDVHPSCKVT
AMKCFLLELQVISLESGDASIHDTCVENLILANNSSLSSNG
NVTESGCCKECEELEEKNIKEFLQSFVHIVQMFINTS

Host: *Escherichia coli*

Theoretical MW (kDa): 12.9

Protocols: See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Form: Lyophilized

Preparation Method: *Escherichia coli* expression system

Purity: > 97% by SDS-PAGE

Activity: ED₅₀ is < 0.5 ng/mL as determined by the dose-dependant stimulation of the proliferation of mouse CTL2-2, cells corresponding to a specific activity of > 2 x 10⁶ units/mg.

Recommend Usage: Biological Activity

SDS-PAGE

The optimal working dilution should be determined by the end user.

Storage Buffer: Lyophilized from sterile distilled Water is > 100 ug/mL

Storage Instruction: Store at 2°C to 8°C for 1 week.

For long term storage, aliquot and store at -20°C to -80°C.

Aliquot to avoid repeated freezing and thawing.

Gene Symbol: IL15

Gene Alias: IL-15, MGC9721

Gene Summary: The protein encoded by this gene is a cytokine that regulates T and natural killer cell activation and proliferation. This cytokine and interleukine 2 share many biological activities. They are found to bind common hematopoietin receptor subunits, and may compete for the same receptor, and thus negatively regulate each other's activity. The number of CD8+ memory cells is shown to be controlled by a balance between this cytokine and IL2. This cytokine induces the activation of JAK kinases, as well as the phosphorylation and activation of transcription activators STAT3, STAT5, and STAT6. Studies of the mouse counterpart suggested that this cytokine may increase the expression of apoptosis inhibitor BCL2L1/BCL-x(L), possibly through the transcription activation activity of STAT6, and thus prevent apoptosis. Two alternatively spliced transcript variants of this gene encoding the same protein have been reported. [provided by RefSeq]