



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## Datasheet

### IL17A (Human) Recombinant Protein

**Catalog Number:** P8358

**Regulation Status:** For research use only (RUO)

**Product Description:** Human IL17A (Q16552, 24 a.a. - 155 a.a.) partial recombinant protein with His tag at C-terminus expressed in Sf9 cells.

**Sequence:**

```
GITIPRNP GCPNSEDKNFPRTVMVNLNIHNRNTNTNPK  
RSSDY YNRSTSPWNLHRNEDPERYPSVIWEAKCRHL  
GCINADGNVDYHMNSVPIQQEILVLRREPPHCPNSFRL  
EKILVSVGCTCVTPIVHHVAHHHHHH
```

**Host:** insect

**Theoretical MW (kDa):** 15.9

**Protocols:** See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

**Form:** Liquid

**Preparation Method:** Sf9 cell expression system

**Purity:** > 90% by SDS-PAGE

**Recommend Usage:** Biological Activity  
SDS-PAGE

The optimal working dilution should be determined by the end user.

**Storage Buffer:** In 20mM PBS pH 7.4 (0.1mM PMSF, 1mM EDTA, 20% glycerol)

**Storage Instruction:** Store at 2°C to 8°C for 1 week.  
For long term storage, aliquot and store at -20°C to -80°C.

Aliquot to avoid repeated freezing and thawing.

**Entrez GeneID:** 3605

**Gene Symbol:** IL17A

**Gene Alias:** CTLA8, IL-17, IL-17A, IL17

**Gene Summary:** The protein encoded by this gene is a

proinflammatory cytokine produced by activated T cells. This cytokine regulates the activities of NF-kappaB and mitogen-activated protein kinases. This cytokine can stimulate the expression of IL6 and cyclooxygenase-2 (PTGS2/COX-2), as well as enhance the production of nitric oxide (NO). High levels of this cytokine are associated with several chronic inflammatory diseases including rheumatoid arthritis, psoriasis and multiple sclerosis. [provided by RefSeq]