



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

IL17B (Human) Recombinant Protein

Catalog Number: P8361

Regulation Status: For research use only (RUO)

Product Description: Human IL17B (Q9UHF5, 21 a.a. - 180 a.a.) partial recombinant protein with His-tag at N-terminus expressed in *Escherichia coli*.

Sequence:

MGSSHHHHHSSGLVPRGSHMGSHMQPRSPKSKRK
GQGRPGPLAPGPHQVPLDLVSRMKPYARMEEYERNI
EEMVAQLRNSSELAQRKCEVNLQLWMSNKRSLSPWG
YSINHDPSPRIPVDLPEARCLCLGCVNPFTMQEDRSMV
SVPVFSQVPVRRRLCPPPPRTGPCRQRAVMETIAVGC
TCIF

Host: *Escherichia coli*

Theoretical MW (kDa): 20

Protocols: See our web site at
<http://www.abnova.com/support/protocols.asp> or product
page for detailed protocols

Form: Liquid

Preparation Method: *Escherichia coli* expression
system

Purity: > 90% by SDS-PAGE

Recommend Usage: Biological Activity
SDS-PAGE

The optimal working dilution should be determined by
the end user.

Storage Buffer: In 20mM Tris-HCl buffer pH 8.0 (0.4M
Urea, 10% glycerol)

Storage Instruction: Store at 2°C to 8°C for 1 week.
For long term storage, aliquot and store at -20°C to
-80°C.
Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 27190

Gene Symbol: IL17B

Gene Alias: IL-17B, IL-20, MGC138900, MGC138901,
ZCYTO7

Gene Summary: The protein encoded by this gene is a
T cell-derived cytokine that shares sequence similarity
with IL17. This cytokine was reported to stimulate the
release of TNF alpha (TNF) and IL1 beta (IL1B) from a
monocytic cell line. Immunohistochemical analysis of
several nerve tissues indicated that this cytokine is
primarily localized to neuronal cell bodies. [provided by
RefSeq]