



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

IL17B (Human) Recombinant Protein

Catalog Number: P8362

Regulation Status: For research use only (RUO)

Product Description: Human IL17B (Q9UHF5, 21 a.a. - 180 a.a.) partial recombinant protein with His-tag at C-terminus expressed in Sf9 cells.

Sequence:

ADPQPRSPKSKRKGQGRPGPLAPGPHQVPLDLVSRM
KPYARMEEYERNIEEMVAQLRNSSELAQRKCEVNLQL
WMSNKRSLSPWGYSINHPSRIPVDLPEARCLCLGCV
NPFTMQEDRSMVSPVFSQVPVRRRLCPPPPRTGPC
RQRAVMETIAVGCTCIFHHHHHH

Host: insect

Theoretical MW (kDa): 19.2

Protocols: See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Form: Liquid

Preparation Method: Sf9 cell expression system

Purity: > 90% by SDS-PAGE

Recommend Usage: Biological Activity

SDS-PAGE

The optimal working dilution should be determined by the end user.

Storage Buffer: In PBS pH 7.4 (10% glycerol)

Storage Instruction: Store at 2°C to 8°C for 1 week.

For long term storage, aliquot and store at -20°C to -80°C.

Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 27190

Gene Symbol: IL17B

Gene Alias: IL-17B, IL-20, MGC138900, MGC138901, ZCYTO7

Gene Summary: The protein encoded by this gene is a T cell-derived cytokine that shares sequence similarity with IL17. This cytokine was reported to stimulate the release of TNF alpha (TNF) and IL1 beta (IL1B) from a monocytic cell line. Immunohistochemical analysis of several nerve tissues indicated that this cytokine is primarily localized to neuronal cell bodies. [provided by RefSeq]