



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

IL17F (Human) Recombinant Protein

Catalog Number: P8367

Regulation Status: For research use only (RUO)

Product Description: Human IL17F (Q96PD4, 31 a.a. - 163 a.a.) partial recombinant protein with His-tag at N-terminus expressed in *Escherichia coli*.

Sequence:

MGSSHHHHHSSGLVPRGSHMGSHMRKIPKVGHTFF
QKPESCPPVPGGSMKLDIGIINENQRVSMRNIESRST
SPWNYVTWDPNRYPSVQVQAQCRNLGCINAQGKED
ISMNSVPIQQETLVVRRKHQGCVSFQLEKVLVTVGCT
CVTPVIHHVQ

Host: *Escherichia coli*

Theoretical MW (kDa): 17.6

Protocols: See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Form: Liquid

Preparation Method: *Escherichia coli* expression system

Purity: > 85% by SDS-PAGE

Recommend Usage: Biological Activity

SDS-PAGE

The optimal working dilution should be determined by the end user.

Storage Buffer: In 20mM Tris-HCl buffer pH 8.0 (0.4M Urea, 10% glycerol)

Storage Instruction: Store at 2°C to 8°C for 1 week.

For long term storage, aliquot and store at -20°C to -80°C.

Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 112744

Gene Symbol: IL17F

Gene Alias: IL-17F, ML-1, ML1

Gene Summary: The protein encoded by this gene is a cytokine that shares sequence similarity with IL17. This cytokine is expressed by activated T cells, and has been shown to stimulate the production of several other cytokines, including IL6, IL8, and CSF2/GM-CSF. This cytokine is also found to inhibit the angiogenesis of endothelial cells and induce endothelial cells to produce IL2, TGFB1/TGFB, and monocyte chemoattractant protein-1. [provided by RefSeq]