



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## Datasheet

### IL17F (Human) Recombinant Protein

**Catalog Number:** P8368

**Regulation Status:** For research use only (RUO)

**Product Description:** Human IL17F (Q96PD4, 31 a.a. - 163 a.a.) partial recombinant protein with His-tag at C-terminus expressed in Sf9 cells.

**Sequence:**

ADPRKIPKVGHTFFQKPESCPPVPGGSMKLDIGIINEN  
QRVSMRNIESRSTSPWNYVTWDPNRYPSEVVQAQ  
CRNLGCINAQGGKEDISMNSVPIQQETLVVRRKHQGCS  
VSFQLEKVLVTVGCTCVTPVIHHVQHHHHHH

**Host:** insect

**Theoretical MW (kDa):** 16

**Protocols:** See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

**Form:** Liquid

**Preparation Method:** Sf9 cell expression system

**Purity:** > 95% by SDS-PAGE

**Recommend Usage:** Biological Activity  
SDS-PAGE

The optimal working dilution should be determined by the end user.

**Storage Buffer:** In PBS pH 7.4 (1mM DTT, 20% glycerol)

**Storage Instruction:** Store at 2°C to 8°C for 1 week.  
For long term storage, aliquot and store at -20°C to -80°C.

Aliquot to avoid repeated freezing and thawing.

**Entrez GeneID:** 112744

**Gene Symbol:** IL17F

**Gene Alias:** IL-17F, ML-1, ML1

**Gene Summary:** The protein encoded by this gene is a

cytokine that shares sequence similarity with IL17. This cytokine is expressed by activated T cells, and has been shown to stimulate the production of several other cytokines, including IL6, IL8, and CSF2/GM-CSF. This cytokine is also found to inhibit the angiogenesis of endothelial cells and induce endothelial cells to produce IL2, TGFB1/TGFB, and monocyte chemoattractant protein-1. [provided by RefSeq]