



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

CSF2 (Human) Recombinant Protein

Catalog Number: P8742

Regulation Status: For research use only (RUO)

Product Description: Human CSF2 partial recombinant protein expressed in insect cells.

Host: insect

Theoretical MW (kDa): 14.6

Protocols: See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Form: Lyophilized

Preparation Method: Insect cell expression system

Purification: chromatographic

Purity: > 98% as determined by SDS-PAGE.

Activity: ED₅₀ < 0.1 ng/mL, measured by the dose-dependant stimulation of the proliferation of human TF-1 cells (human erythroleukemic indicator cell line), corresponding to a specific activity of 1.0 x 10⁷ IU/mg.

Storage Buffer: Protein (1 mg/mL) was lyophilized from PBS. Reconstitute the lyophilized powder in ddH₂O to 100 ug/mL.

Storage Instruction: Lyophilized protein at room temperature for 3 weeks, should be stored at -20°C. Protein aliquots at 4°C for 2-7 days and should be stored at -20°C to -80°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).
Avoid repeated freeze/thaw cycles.

Entrez GeneID: 1437

Gene Symbol: CSF2

Gene Alias: GMCSF, MGC131935, MGC138897

Gene Summary: The protein encoded by this gene is a cytokine that controls the production, differentiation, and

function of granulocytes and macrophages. The active form of the protein is found extracellularly as a homodimer. This gene has been localized to a cluster of related genes at chromosome region 5q31, which is known to be associated with interstitial deletions in the 5q- syndrome and acute myelogenous leukemia. Other genes in the cluster include those encoding interleukins 4, 5, and 13. [provided by RefSeq]