



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

PDGFB (Human) Recombinant Protein

Catalog Number: P9010

Regulation Status: For research use only (RUO)

Product Description: Human PDGFB (P01127, 82 a.a. - 190 a.a.) partial-length recombinant protein with His tag at N-Terminus expressed in *Escherichia coli*.

Host: *Escherichia coli*

Theoretical MW (kDa): 16.75

Protocols: See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Form: Liquid

Preparation Method: *Escherichia coli* expression system

Purity: > 95% by SDS PAGE

Storage Buffer: 25mM Na-Acetate pH 4.8, 1mM EDTA and 50% glycerol.

Storage Instruction: Store at -20°C. Aliquot the product after reconstitution to avoid repeated freezing/thawing cycles.

Entrez GeneID: 5155

Gene Symbol: PDGFB

Gene Alias: FLJ12858, PDGF2, SIS, SSV, c-sis

Gene Summary: The protein encoded by this gene is a member of the platelet-derived growth factor family. The four members of this family are mitogenic factors for cells of mesenchymal origin and are characterized by a motif of eight cysteines. This gene product can exist either as a homodimer (PDGF-BB) or as a heterodimer with the platelet-derived growth factor alpha polypeptide (PDGF-AB), where the dimers are connected by disulfide bonds. Mutations in this gene are associated with meningioma. Reciprocal translocations between chromosomes 22 and 7, at sites where this gene and

that for COL1A1 are located, are associated with a particular type of skin tumor called dermatofibrosarcoma protuberans resulting from unregulated expression of growth factor. Two alternatively spliced transcript variants encoding different isoforms have been identified for this gene. [provided by RefSeq]