



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## Datasheet

### CD84 (Human) Recombinant Protein

**Catalog Number:** P9110

**Regulation Status:** For research use only (RUO)

**Product Description:** Human CD84 (Q9UIB8, 22 a.a. - 225 a.a.) partial-length recombinant protein with His tag at N-Terminus expressed in *Escherichia coli*.

**Sequence:**

MGSSHHHHHSSGLVPRGSHMGSHMKDSEIFTVNGI  
LGESVTFPVNIQEPQVKIIAWTSKTSVAYVTPGDSET  
APVVTVTHRNYERIHALGPNYNLVISDLRMEDAGDYK  
ADINTQADPYTTTKRYNLQIYRRLGKPKITQSLMASVN  
STCNVTLTCSVEKEEKNVTYNWSPLGEEGNVLQIFQT  
PEDQELTYTCTAQNPVSNNSDSISARQLCADIAMGFR  
THHTG.

**Host:** *Escherichia coli*

**Theoretical MW (kDa):** 25.4

**Protocols:** See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

**Form:** Liquid

**Preparation Method:** *Escherichia coli* expression system

**Purity:** > 80% by SDS-PAGE.

**Storage Buffer:** 20mM Tris-HCl buffer (pH 8.0), 0.4M urea and 10% glycerol.

**Storage Instruction:** Store at -20°C. Aliquot the product after reconstitution to avoid repeated freezing/thawing cycles.

**Entrez GeneID:** 8832

**Gene Symbol:** CD84

**Gene Alias:** DKFZp781E2378, LY9B, SLAMF5, hCD84, mCD84

**Gene Summary:** Members of the CD2 (see MIM 186990) subgroup of the Ig superfamily, such as CD84,

have similar patterns of conserved disulfide bonds and function in adhesion interactions between T lymphocytes and accessory cells.[supplied by OMIM]