



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

TNFRSF13B (Human) Recombinant Protein

Catalog Number: P9190

Regulation Status: For research use only (RUO)

Product Description: Human TNFRSF13B full-length recombinant protein with His tag in N-terminus expressed in *Escherichia coli*.

Sequence:

MGSSHHHHHSSGLVPRGSHMGSMGLGRSRRGGR
SRVDQEERFPQGLWTGVAMRSCPEEQYWDPLLGTG
MSCKTICNHQSQRRTCAAFCSRSLSCRKEQGKFDHLLR
DCISCASICGQHPKQCAYFCENKLRSPVNLPELRRQ
RSGEVENNSDNSGRYQGLEHRGSEASPALPGLKLSA
DQVALVYS

Host: *Escherichia coli*

Theoretical MW (kDa): 20.9

Protocols: See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Specificity: TACI Human, His

Form: Liquid

Preparation Method: *Escherichia coli* expression system

Purification: chromatographic

Purity: > 90% as determined by SDS-PAGE.

Storage Buffer: Solution (1 mg/mL) containing 20 mM Tris-HCl, pH 8.0, 10% glycerol, 0.4 M Urea.

Storage Instruction: Store at 4°C for 2-4 weeks and should be stored at -20°C to -80°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).
Avoid repeated freeze/thaw cycles.

Entrez GeneID: 23495

Gene Symbol: TNFRSF13B

Gene Alias: CD267, CVID, FLJ39942, MGC133214, MGC39952, TACI, TNFRSF14B

Gene Summary: The protein encoded by this gene is a lymphocyte-specific member of the tumor necrosis factor (TNF) receptor superfamily. It interacts with calcium-modulator and cyclophilin ligand (CAML). The protein induces activation of the transcription factors NFAT, AP1, and NF-kappa-B and plays a crucial role in humoral immunity by interacting with a TNF ligand. This gene is located within the Smith-Magenis syndrome region on chromosome 17. [provided by RefSeq]