



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## Datasheet

### TNFRSF10A (Human) Recombinant Protein

TRAILR1

**Catalog Number:** P9260**Regulation Status:** For research use only (RUO)**Product Description:** Human TNFRSF10A partial recombinant protein with His tag in C-terminus expressed in Baculovirus cells.**Sequence:**

```
ASGTEAAAATPSKVWGSSAGRIEPRGGGRGALPTSM
GQHGPSARARAGRAPGPRPAREASPLRVHKTFFKVV
VVGVLQVVPSSAATIKLHDQSIGTQQWEHSPLGELCP
PGSHRSEHPGACNRCTEGVGYTNASNLFACLPTCA
CKSDEEERSPCTTTRNTACQCKPGTFRNDNSAEMCR
KCSRGCPRGMVKVKDCTPWSDIECVHKESGNGHNLE
HHHHHH
```

**Host:** Viruses**Theoretical MW (kDa):** 23.9**Protocols:** See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols**Form:** Liquid**Preparation Method:** Baculovirus expression system**Purification:** chromatographic**Purity:** > 90% as determined by SDS-PAGE.**Storage Buffer:** Solution (0.5 mg/mL) containing 1X PBS, pH 7.4, 10% glycerol.**Storage Instruction:** Store at 4°C for one weeks and should be stored at -20°C to -80°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).  
Avoid repeated freeze/thaw cycles.**Entrez GeneID:** 8797**Gene Symbol:** TNFRSF10A**Gene Alias:** APO2, CD261, DR4, MGC9365, TRAILR-1,**Gene Summary:** The protein encoded by this gene is a member of the TNF-receptor superfamily. This receptor is activated by tumor necrosis factor-related apoptosis inducing ligand (TNFSF10/TRAIL), and thus transduces cell death signal and induces cell apoptosis. Studies with FADD-deficient mice suggested that FADD, a death domain containing adaptor protein, is required for the apoptosis mediated by this protein. [provided by RefSeq]